


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CG/1 Lucerne - Single and Repeated Applications of K

Rothamsted Research

Rothamsted Research (1957) *56/R/CG/1 Lucerne - Single and Repeated Applications of K* ; Yields Of The Field Experiments 1956, pp 111 - 112 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Cg/1.1

LUCERNE

Single and repeated applications of potash - Great Harpenden II 1956, the second year.

Design: 6 randomized blocks of 8 plots each.

Area of each plot: 0.0147 acres.

Treatments: (For the 3 years 1955 - 57)

None (2 plots per block).

Single dressings in 1955: 1; 2; 3 cwt K_2O per acre as muriate of potash.

Annual dressings: 0.33; 0.66; 1.0 cwt K_2O per acre as muriate of potash.

Note: In spring 1955 all the potash dressings were given in the seed bed. Subsequent annual applications are applied as top dressings.

Basal dressing 1956: 3 tons ground chalk per acre.

Cultivations, etc.: Annual potash dressings applied: Nov 8, 1955.

Ground chalk applied: Mar 12, 1956. Cut and weighed green: three times - June 11, July 24, Oct 24.

Standard errors per plot. Dry matter:

1st cut:	2.15 cwt per acre or 7.4% (36 d.f.)
2nd cut:	2.50 cwt per acre or 9.4% (36 d.f.)
3rd cut:	0.972 cwt per acre or 5.8% (36 d.f.)
Total of 3 cuts:	3.89 cwt per acre or 5.4% (36 d.f.)

Note: For the first year's results, see "Results of the Field Experiments" 1955, Section Ce/1.

56/Cg/1.2

Summary of Results

Dry matter: cwt per acre								
K ₂ O: cwt per acre								
Applied annually								
Applied 1955 1955 and 1956								
None	1.0	2.0	3.0	0.33	0.66	1.0	Mean	
1st cut								
	(± 0.62)			(± 0.88)				
Mean	27.2	29.7	30.8	31.9	27.6	29.8	29.8	29.3
Increase (±1.08)		2.5	3.6	4.7	0.4	2.6	2.6	
2nd cut								
	(± 0.72)			(± 1.02)				
Mean	25.5	26.0	28.1	27.6	25.9	26.7	27.1	26.5
Increase (±1.25)		0.5	2.6	2.1	0.4	1.2	1.6	
3rd cut								
	(± 0.28)			(± 0.40)				
Mean	16.0	16.8	17.6	17.3	16.4	17.1	17.3	16.8
Increase (±0.49)		0.8	1.6	1.3	0.4	1.1	1.3	
Total of 3 cuts								
	(± 1.12)			(± 1.59)				
Mean	68.7	72.5	76.5	76.8	69.8	73.6	74.2	72.6
Increase (±1.95)		3.8	7.8	8.1	1.1	4.9	5.5	
1st cut, Mean dry matter % as cut:								
								24.2
2nd cut, Mean dry matter % as cut:								
								23.0
3rd cut, Mean dry matter % as cut:								
								24.5
Total of 3 cuts, Mean dry matter % as cut:								
								23.9