


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1956

[Full Table of Content](#)



---

### 56/R/BA/3 and 56/W/BA/3 Six-course Rotation

#### Rothamsted Research

Rothamsted Research (1957) *56/R/BA/3 and 56/W/BA/3 Six-course Rotation* ; Yields Of The Field Experiments 1956, pp 25 - 28 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Ba/3.1

SIX COURSE ROTATION EXPERIMENT

The 27th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1956.

For history, treatments etc., see "Details of the Classical and Long Term Experiments", 1956.

Area of each plot: Rothamsted, 0.0250 acres. Woburn, 0.0266 acres.

Cultivations, etc.:

Rothamsted

Sugar beet.

Ploughed: Sept 6, 1955 and Nov 24. Fertilizers applied, seed drilled at 18 lb per acre: Mar 29, 1956. Singled: May 24 - 30. Lifted: Dec 5. Variety: Klein E.

Barley.

Sugar beet tops spread: Nov 22, 1955. Ploughed: Nov 24. Ground chalk applied at 20 cwt per acre: Dec 2. Seed drilled at 3 bushels per acre: Mar 14, 1956. Fertilizers applied: Mar 15. Clover seed undersown: Apr 23. Harvested: Sept 12. Variety: Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 26, 1955. Autumn fertilizers applied: Oct 27. Sulphate of ammonia applied: Apr 9, 1956. Cut: July 13. Variety: Late flowering Montgomery Red.

Wheat.

Ploughed: July 28, 1955 and Oct 12. Autumn fertilizers applied: Oct 17. Seed drilled at  $2\frac{1}{4}$  bushels per acre: Oct 18. Sulphate of ammonia applied: Apr 24, 1956. Harvested: Aug 22. Variety: Yeoman.

Potatoes.

Ploughed: Sept 6, 1955 and Oct 17. Ridged: Mar 26, 1956. Fertilizers applied, potatoes planted: Mar 27. Earthed up: June 23. Sprayed with copper fungicide at 3 lb in 80 gallons per acre: July 25. Sprayed with sulphuric acid, 20% BOV, 80 gallons per acre: Sept 13. Lifted: Oct 8. Variety: Majestic.

Rye.

Ploughed: Oct 3, 1955. Ground chalk applied at 20 cwt per acre: Oct 4. Autumn fertilizers applied: Oct 15. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 24, 1956. Harvested: Aug 21. Variety: King II.



56/Ba/3.2

Woburn

Sugar beet.

Ploughed: Sept 1, 1955 and Dec 12. Fertilizers applied: Apr 6, 1956.  
Seed drilled at 12 lb per acre: Apr 9. Re-drilled thin places:  
May 12. Dusted with 5% DDT: May 19. Sprayed with parathion,  
 $\frac{1}{2}$  pint in 40 gallons per acre: May 25. Singled: June 6 - 7.  
Lifted: Oct 23. Variety: Klein E.  
Note: The DDT and parathion were applied to control leaf miner  
(Pegomyia beta).

Barley.

Beet tops spread, ploughed: Oct 31, 1955. Fertilizers applied:  
Mar 13. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Mar 15. Sprayed  
with MCPA, 2 pints in 20 gallons per acre: May 7. Harvested:  
Aug 14. Variety: Herta.

Clover.

Autumn fertilizers applied: Oct 13, 1955. Undersown crop failed  
during winter. Whole block ploughed: Mar 9, 1956. Sulphate of  
ammonia applied: Mar 27. Seed drilled at 40 lb per acre: Mar 28.  
Cut: July 6. Variety: Crimson Clover.

Wheat.

Ploughed: July 19, 1955. Autumn fertilizers applied: Oct 12, 1955.  
Seed drilled at 3 bushels per acre: Oct 15. Sulphate of ammonia  
applied: Apr 26, 1956. Sprayed with DNOC, 6 lb in 80 gallons  
per acre: May 1. Harvested: Aug 23. Variety: Yeoman.

Potatoes.

Ploughed: Sept 1, 1955 and Dec 12. Ridged, fertilizers applied,  
potatoes planted: Apr 4, 1956. Earthed up: June 18. Sprayed with  
copper fungicide, 5 lb in 40 gallons per acre: July 23. Sprayed with  
arsenious compound, 1 gallon in 40 gallons per acre: Sept 4.  
Lifted: Oct 4. Variety: Majestic.

Rye.

Ploughed: Oct 6, 1955. Ground chalk applied at 20 cwt per acre:  
Oct 13. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Oct 15. Seed  
re-drilled at  $2\frac{1}{2}$  bushels per acre: Nov 16. Sulphate of ammonia  
applied: Apr 24, 1956. Harvested: Aug 23. Variety: King II.  
Note: A poor plant was obtained from the first sowing because of  
a defective drill.

Note: In 1956 at Woburn the nitrogen levels were doubled on all crops  
except clover.



56/Ba/3.3

Summary of Results

Mean yields per acre and responses in yield per cwt of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O

	Rothamsted	Woburn	Rothamsted	Woburn
Sugar Beet, roots (washed): tons per acre			Barley, grain: cwt per acre	
Mean	11.59	10.76	*	
Response to: N	+5.27	+8.97	25.3	34.0
P	-2.55	+2.21	+16.9	+23.6
K	+0.07	+0.64	-0.3	-1.0
Mean dry matter % as harvested:			-3.7	-2.7
			80.9	
Sugar Beet, sugar percentage			Barley, straw: cwt per acre	
Mean	18.4	18.1	*	
Response to: N	-0.4	0.0	26.9	29.2
P	0.0	-0.1	+26.3	+25.1
K	+0.1	+0.2	+0.5	-1.2
Mean dry matter % as harvested:			-6.4	-2.3
			81.4	
Sugar Beet, total sugar: cwt per acre			Clover, hay, dry matter: cwt per acre	
Mean	42.6	38.9	21.3	19.1
Response to: N	+18.5	+32.1	+5.3	-5.5
P	-9.1	+7.7	-11.7	+12.4
K	+0.6	+2.6	-2.5	+12.9
Mean dry matter % as harvested:			79.7	
Sugar Beet, tops: tons per acre			Wheat, grain: cwt per acre	
Mean	9.22	6.41	*	
Response to: N	+6.32	+8.08	27.7	19.1
P	-4.37	+2.65	+6.1	+11.3
K	-0.31	+0.46	+5.7	-1.0
Mean dry matter % as harvested:			+1.6	-0.2
			78.1	
Sugar Beet, plant number: thousands per acre			Wheat, straw: cwt per acre	
Mean	28.1	**	*	
Response to: N	-2.1		37.6	23.2
P	-3.7		+14.5	+10.9
K	+1.1		+10.5	-2.5
Mean dry matter % as harvested:			-0.6	-1.2
			79.9	

\*(at 85% dry matter)

\*\* not recorded.



56/Ba/3.4

Mean yields per acre and responses in yield per cwt of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O

	Rothamsted	Woburn	Rothamsted	Woburn
	Potatoes, total tubers: tons per acre		Rye, grain: cwt per acre	
			*	
Mean	9.03	10.87	28.8	30.7
Response to: N	+6.19	+5.33	+19.1	+15.5
P	+1.83	+2.61	+2.0	+1.9
K	+0.42	-0.04	-4.7	-5.7
Mean dry matter % as harvested:			76.6	
	Potatoes, percentage ware		Rye, straw: cwt per acre	
	(1)	(2)	*	
Mean	91.0	83.3	36.4	35.1
Response to: N	+7.3	+6.2	+23.1	+17.6
P	-17.1	-13.8	+2.3	+0.6
K	+0.2	-9.8	-6.9	-6.4
Mean dry matter % as harvested:			80.7	

\*(at 85% dry matter)

Riddle: (1) 1½"; (2) 1⅝".