


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/A/3 Agdell - Winter Beans

Rothamsted Research

Rothamsted Research (1957) *56/R/A/3 Agdell - Winter Beans* ; Yields Of The Field Experiments 1956, pp 7 - 7 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1956

Without manure 1851 and since

For history, treatments etc. see "Details of the Classical and Long Term Experiments" 1956.

Cultivations, etc.:

Cropped plots. Ploughed: Sept 28, 1955. Seed drilled at 3 bushels per acre: Nov 2. Harvested: Sept 1, 1956. Variety: Squareheads Master 13/4.

Fallowed plots. Ploughed: Sept 28, 1955, May 15, 1956 and Aug 23.

Summary of Results

Mean yields (at 85% dry matter): cwt per acre

Plot	A2	A3	A4	
No. of years of fallow	1	1	3	Mean
Grain	10.2	11.3	10.7	10.7
Straw	15.3	17.1	16.0	16.1

Mean dry matter % as threshed, Grain: 80.5
Straw: 81.9

WINTER BEANS - AGDELL 1956

For history, treatments etc. see "Details of the Classical and Long Term Experiments" 1956. In 1956 a crop of winter sown beans was grown without manure.

Area of plot harvested: 0.0458 acres.

Cultivations, etc.:

Ploughed: Oct 8 - 10, 1955. Seed drilled at 275 lbs per acre: Oct 29. Combine harvested: Oct 4 - 6. Variety: S.Q.Giant.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Manure to turnips 1948	Mineral manure				Complete mineral and nitrogenous manure		Mean
	None since 1848		No nitrogen				
	5	6	3	4	1	2	
Plot rotation	Fallow	Clover	Fallow	Clover	Fallow	Clover	
	8.8	5.2	26.2	20.0	18.2	19.2	16.3

Mean dry matter % as harvested; 73.7