


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CA/3 Winter Wheat - Incidence of Bulb Fly (Seed Rates)

Rothamsted Research

Rothamsted Research (1957) *56/R/CA/3 Winter Wheat - Incidence of Bulb Fly (Seed Rates)* ; Yields Of The Field Experiments 1956, pp 73 - 74 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Ca/3.1

WINTER WHEAT

Seed rates in relation to control of wheat bulb fly - Long Hoos VII
the 3rd (final) year.

Design: 4 4x4 squares with treatments on rows in 1954 and on columns
in 1955.

Area of each plot: 0.0151 acres. Area harvested: 0.0050 acres.

Treatments to spring wheat 1954 and to winter wheat 1955.

Bare fallow.

Seed rates: $\frac{1}{3}$; 1; 3 bushels per acre.

Basal dressings per acre for 1956 crop: 1 cwt compound granular
fertilizer (12% N, 12% P₂O₅, 15% K₂O) combine drilled with seed;
0.8 cwt N as "Nitro-Chalk".

Cultivations, etc.:

Floughed: Sept 14, 1955. Seed combine drilled (all at $2\frac{3}{4}$ bushels
per acre): Oct 28. 'Nitro-Chalk' applied: Apr 30, 1956.

Combine harvested: Aug 31. Variety: Cappelle.

Note: seed treated with organo-mercurial fungicide only.

Standard errors per plot. Grain (at 85% dry matter):

Row: 1.94 cwt per acre or 5.7% (9 d.f.)

Column: 2.65 cwt per acre or 7.8% (9 d.f.)

Plot: 2.03 cwt per acre or 6.0% (27 d.f.)

Records were made of:

Number of wheat bulb fly larvae, weight per ear, number of grains
per ear and plant number.

56/Ga/3.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

1954 Treatments		1955 Treatments				Mean
		Fallow	Seed rate: bushels per acre			
			$\frac{1}{3}$	1	3	
			(1)	(2)	(3) and (4)	(± 0.97)
Seed rate:	Fallow	50.4	37.1	37.7	36.2	40.4
	$\frac{1}{3}$	47.7	30.8	28.4	23.3	32.6
bushels	1	47.3	28.1	27.8	20.5	30.9
per acre	3	47.5	31.1	26.6	23.1	32.1
Mean (± 1.32)		48.2	31.8	30.1	25.8	34.0

- (1) ± 1.59 for use in horizontal comparisons only.
- (2) ± 1.31 for use in vertical comparisons only.
- (3) ± 1.79 for use in diagonal comparisons only.
- (4) ± 1.01 for use in interaction comparisons only.

Mean dry matter % as harvested: 74.9