


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/W/CA/4 Winter Wheat - Varieties, Seed Rates, Levels and Times of N

Rothamsted Research

Rothamsted Research (1957) *56/W/CA/4 Winter Wheat - Varieties, Seed Rates, Levels and Times of N*; Yields Of The Field Experiments 1956, pp 75 - 76 - DOI:

<https://doi.org/10.23637/ERADOC-1-176>

56/Ca/4.1

WINTER WHEAT

Varieties, seed rates, levels and times of application of N - Woburn, Roadpiece 1956, the 3rd year.

Design: 4 randomized blocks of 8 plots each, certain high order interactions being confounded with block differences. In addition each block contained 2 plots with no nitrogen, the variety \times seed rate interaction being confounded.

Area of each plot: 0.0182 acres. Area harvested: 0.0138 acres.

Treatments: All combinations of:-

Varieties: Holdfast; Cappelle.

Seed rates: Holdfast, $1\frac{1}{2}$; 3 bushels per acre.
Cappelle, 2; 4 bushels per acre.

Nitrogen: 0.5; 1.0 cwt N per acre applied as 'Nitro-Chalk'.

Time of application of N: half dressing in March and again in May;
whole dressing mid March; mid April; mid May.

Basal dressing: 1 cwt per acre compound granular fertilizer (12% N, 12% P₂O₅, 15% K₂O) combine drilled with seed.

Cultivations, etc.: Ploughed: Sept 29, 1955. Combine drilled: Oct 26.
March top dressing applied: Mar 8, 1956. April top dressing applied: Apr 12. All plots sprayed with DNOC at 8 lb in 90 gallons per acre: May 1. May top dressing applied: May 17. Combine harvested: Sept 13. Previous crop: Wheat.

Standard error per plot.

Grain (at 85% dry matter): 1.82 cwt per acre or 17.6% (12 d.f.)

Note (1) The experiment is a repetition on the same plots of the ones carried out in 1955 and 1954 (see "Results of the Field Experiments" 55/Ca/4 and 54/Ca/7.)

(2) The crop was severely and irregularly infested with weeds particularly twitch (*Agrostis gigantea*) and Mayweed (*Matricaria*).

(3) Records of incidence of disease (Take-all and Eyespot) and weeds and counts of plant, shoot and ear numbers were made.

56/Ca/4.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	T ₁	T ₂	T ₃	T ₄	Mean	
Mean (±0.65)	12.1	12.8	14.7	6.9	11.6	
	(±0.91)					
V ₁	10.7	10.2	12.8	6.8	10.1	
V ₂	13.5	15.5	16.5	6.9	13.1	
Difference (±1.29)	+2.8	+5.3	+3.7	+0.1	+3.0 (±0.65)	
R ₁	10.1	12.0	11.9	4.7	9.7	
R ₂	14.0	13.7	17.5	9.0	13.6	
Difference (±1.29)	+3.9	+1.7	+5.6	+4.3	+3.9 (±0.65)	
N ₁	11.2	10.9	10.6	7.8	10.1	
N ₂	12.9	14.8	18.7	5.9	13.1	
Difference (±1.29)	+1.7	+3.9	+8.1	-1.9	+3.0 (±0.65)	
	R ₁	R ₂	N ₀	N ₁	N ₂	Mean
Mean			(±0.65)	(±0.46)		
			5.3	10.1	13.1	10.3
	(±0.65)		(±0.91)	(±0.65)		(±0.41)
V ₁	9.0	11.2	5.1	8.7	11.6	9.1
V ₂	10.3	15.9	5.5	11.6	14.6	11.6
R ₁			5.3	8.4	10.9	8.8
R ₂			5.3	11.8	15.3	11.9

Mean dry matter % as harvested: 78.7

Treatments

V₁ Holdfast
V₂ Cappelle

R₁, R₂ 1½, 3 bushels per acre
R₁, R₂ 2, 4 bushels per acre

N₀ No N
N₁ 0.46 cwt N per acre
N₂ 0.93 cwt N per acre

T₁ 'Nitro-Chalk' half in March half in May
T₂ 'Nitro-Chalk' all in mid March
T₃ 'Nitro-Chalk' all in mid April
T₄ 'Nitro-Chalk' all in mid May

The V × R table does not include the plots receiving no nitrogen.