


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1956

[Full Table of Content](#)



---

### 56/R/CD/2 Spring Beans - Control of Weeds (Spraying and Cultivations)

#### Rothamsted Research

Rothamsted Research (1957) *56/R/CD/2 Spring Beans - Control of Weeds (Spraying and Cultivations)* ; Yields Of The Field Experiments 1956, pp 84 - 85 - DOI:  
<https://doi.org/10.23637/ERADOC-1-176>

56/Ca/2.1

### SPRING BEANS

Control of weeds by spraying and cultivations - Great Field 1 1956.

Design: 4 randomized blocks of 7 plots each, 2 blocks being sprayed and 2 unsprayed.

Area of each plot: 0.0194 acres. Area harvested: 0.0139 acres.

Treatments: All combinations of:-

Spraying (on blocks): None; DNEP 5 pints in 80 gallons per acre.

Treatment cultivations:

	<u>Additional cultivations</u>	
With inter-row cultivations	None	(1)
	Harrowed once	(2)
	Harrowed three times	(3)
	Mechanical weeder once	(4)
	Mechanical weeder three times	(5)
Without inter-row cultivations	Harrowed as required	(6)
	Mechanical weeder as required	(7)

Basal dressing: 6 cwt compound granular fertilizer (10% P<sub>2</sub>O<sub>5</sub>, 20% K<sub>2</sub>O) per acre.

Cultivations, etc.: Ploughed: Nov 15, 1955. Basal fertilizer applied: Mar 13, 1956. Seed drilled at 200 lb per acre: Mar 14.

Appropriate blocks sprayed with DNEP, 5 pints in 80 gallons per acre: May 7. Inter-row cultivated: Treatments (2), (3), (4), (5) - May 23.

(1) - June 6. Combine harvested: Oct 12. Variety: Spring Tick.

Previous crop: Barley.

Additional treatment cultivations:

(2) and (4): Apr 27.

(3), (5) and (6): Apr 27, May 4, May 9.

(7): Apr 27, May 4 (twice), May 9.

Standard error per plot:

Grain (at 85% dry matter): 1.52 cwt per acre or 7.1% (12 d.f.)



56/cd/2.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Spray	Treatment cultivations							Mean
	1	2	3	4	5	6	7	
	(±1.07)*							
None	21.3	20.6	18.8	17.9	20.7	21.1	19.1	19.9
DNBP	20.6	23.2	23.2	25.0	22.7	22.5	21.7	22.7
Mean (±0.76)	21.0	21.9	21.0	21.4	21.7	21.8	20.4	21.3
Diff. (±1.52) <sup>+</sup>	-0.7	+2.6	+4.4	+7.1	+2.0	+1.4	+2.6	+2.8

\*for use in horizontal comparisons only.

<sup>+</sup>for use in the comparison of two differences only.

Mean dry matter % as harvested: 73.9

Treatment cultivations:

- |                                   |                                |
|-----------------------------------|--------------------------------|
| (1) None                          | With inter-row cultivations    |
| (2) Harrowed once                 | " " " " "                      |
| (3) Harrowed three times          | " " " " "                      |
| (4) Mechanical weeder once        | " " " " "                      |
| (5) Mechanical weeder three times | " " " " "                      |
| (6) Harrowed as required          | Without inter-row cultivations |
| (7) Mechanical weeder as required | " " " " " "                    |