


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CD/3 Spring Beans - Flower Drop (Hormone Sprays)

Rothamsted Research

Rothamsted Research (1957) *56/R/CD/3 Spring Beans - Flower Drop (Hormone Sprays)* ; Yields Of The Field Experiments 1956, pp 86 - 86 - DOI: <https://doi.org/10.23637/ERADOC-1-176>

56/Oa/3

SPRING BEANS

Flower drop - hormone sprays - Great Field I 1956.

Design: 4 randomized blocks of 6 plots each.

Area of each plot: 0.0064 acres. Area harvested: 0.0046 acres.

Treatments: Hormone sprays

- None (2 plots per block) (0)
- 2; 4 applications of 4-chlorophenoxyacetic acid (1) & (2)
- 2; 4 applications of α (2, 4, 5 trichlorophenoxy) propionic acid (3) & (4)

The sprays were applied at a concentration of 5 p.p.m.

Basal dressing: 6 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Cultivations, etc.: Ploughed: Nov 10 - 15, 1955. Basal fertilizer applied: Mar 13, 1956. Seed drilled at 200 lb per acre: Mar 14. All plots (except control) sprayed with hormone sprays at approx. 150 gallons followed by 250 gallons per acre: June 18 and June 25. Appropriate plots sprayed again with hormone sprays at 350 gallons per acre: July 4 and July 10. Combine harvested: Oct 12. Variety: Spring Tick. Previous crop: Barley.

Standard error per plot:

Grain (at 85% dry matter): 1.61 cwt per acre or 6.6% (16 d.f.)

Note: Counts of numbers of pods were made.

Summary of Results

Grain (at 85% dry matter): cwt per acre

		Treatment					
		0	1	2	3	4	Mean
Mean	(±0.80)	24.4 ⁽¹⁾	25.7	27.0	23.0	22.7	24.5
Increase	(±0.99)		+1.3	+2.6	-1.4	-1.7	

(1) ±0.57

Mean dry matter % as harvested: 72.7