


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CD/4 Beans and 56/W/CD/4 - Autumn and Spring Sowing , Spraying, P and K

Rothamsted Research

Rothamsted Research (1957) *56/R/CD/4 Beans and 56/W/CD/4 - Autumn and Spring Sowing ,
Spraying, P and K* ; Yields Of The Field Experiments 1956, pp 87 - 89 - DOI:
<https://doi.org/10.23637/ERADOC-1-176>

56/Ca/4.1

BEANS

Time of sowing, spraying, P and K - Rothamsted (R) Great Harpenden II and Woburn (W) Broadmead I 1956.

Design: 3 blocks of 4 whole plots each split into 3, with spraying on pairs of whole-plots and PK partially confounded.

Area of each sub-plot: Rothamsted, 0.0283 acres; Woburn, 0.0337 acres.
Area harvested: Rothamsted, 0.0088 acres; Woburn, 0.0105 acres.

Treatments. All combinations of:

Time of sowing: Autumn; spring.

Spray: None; "Metasystox" at 2 pints in 80 gallons per acre.

Phosphate: None; 0.5; 1.0 cwt P_2O_5 per acre as superphosphate.

Potash: None; 1.0; 2.0 cwt K_2O per acre as muriate of potash.

Basal dressing: None.

Note. At Woburn the autumn sown beans failed because of bird damage.

Cultivations, etc.:

Rothamsted. Ploughed: Oct 3, 1955. Fertilizers applied for autumn beans, seed drilled at 300 lb per acre: Oct 18. Fertilizers applied for spring beans: Mar 7, 1956. Spring beans sown at 200 lb per acre: Mar 10. Appropriate plots sprayed: June 23. Combine harvested: Oct 4. Previous crop: Barley.

Woburn. Ploughed: Oct 21, 1955. Fertilizers applied for winter beans: Nov 1. Ground chalk at 18 cwt per acre applied: Nov 14. Seed drilled at 275 lb per acre: Nov 15. Fertilizers applied for spring beans: Mar 20, 1956. Seed drilled at 200 lb per acre: Mar 22. Appropriate plots sprayed: June 26. Combine harvested: Oct 23. Variety: Winter-S.Q.Giant, Spring-Albyn.

Standard errors per plot. Grain (at 85% dry matter): cwt per acre.

Great Harpenden II (R), whole plot: 3.06 cwt per acre or 17.3%
(4 d.f.)

sub plot: 2.59 cwt per acre or 14.6%
(12 d.f.)

Broadmead I (W),

Spring beans sub plot: 2.28 cwt per acre or 13.7%
(4 d.f.)

Summary of Results

Great Harpenden II (Rothamsted)

Grain (at 85% dry matter): cwt per acre

	Sown		Diff- erence	F205: cwt per acre		K20: cwt per acre		Mean
	Autumn	Spring		None	1.0	None	2.0	
<u>Spray</u>								
None	17.9	16.1	-1.8	17.5	17.6	15.7	17.6	17.0
Metasystox	19.0	17.8	-1.2	17.9	18.8	15.6	20.6	18.4
Difference	+1.1	+1.7	+0.6	+0.4	+1.2	-0.1	+3.0	+1.4
			(± 3.53)		(± 1.06)*		(± 1.06)*	
					(± 1.49)**		(± 1.49)**	
<u>Sown</u>					(1) & (2)		(1) & (2)	
Autumn				17.8	19.6	15.0	20.4	18.4
Spring				17.6	16.8	16.3	17.8	17.0
Mean				17.7	18.2	15.7	19.1	17.7
Difference				-0.2	-2.8	+1.3	-2.6	-1.4
(± 2.15)				(± 0.75)	-1.3		(± 0.75)	(± 1.77)

Mean dry matter % as harvested: 72.2

*For use in horizontal comparisons only

**For use only in testing the difference of two differences

(1) ±1.52 for use in diagonal comparisons only

(2) ±1.06 for use in horizontal comparisons only

Broadmead I (Woburn)

56/Oa/4.3

Spring beans

Grain (at 85% dry matter): cwt per acre

Spray	P ₂ O ₅ : cwt per acre			K ₂ O: cwt per acre			Mean	
	None	0.5	1.0	None	1.0	2.0		
	(± 1.32)*			(± 1.32)*				
None	15.4	15.7	15.0	10.9	15.4	19.9	15.4	
Metasystox	20.6	17.5	15.6	14.0	17.0	22.7	17.9	
Mean	(±0.93)	18.0	16.6	15.3	12.5	16.2	21.3	16.7
Difference	(±1.86)**	5.2	1.8	0.6	3.1	1.6	2.8	2.5

*For use in horizontal comparisons only

**For use only in testing the difference of 2 differences

Mean dry matter % as harvested: 66.5