


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1956

[Full Table of Content](#)



56/R/CE/5 and 56/W/CE/5 Potatoes - Control of Skin Spot (Fungicides)

Rothamsted Research

Rothamsted Research (1957) *56/R/CE/5 and 56/W/CE/5 Potatoes - Control of Skin Spot (Fungicides)* ;
Yields Of The Field Experiments 1956, pp 99 - 100 - DOI:
<https://doi.org/10.23637/ERADOC-1-176>

56/Ce/5.1

POTATOES

Control of skin spot (Oospora pustulans) by fungicides - Great Knott
III 1956.

Design: 3 randomized blocks of 15 plots each.

Area of each plot: 0.0140 acres. Area harvested: 0.0070 acres.

Treatments:

None (3 plots per block).

PCNB, 20% Parachloronitro benzene dust applied: to seed at 3 oz.
per cwt (S); to furrows before planting at 300 lb per acre (F);
(S) and (F).

Calomel, 4% dust applied: to seed at 12 oz. per cwt (S); to
furrows before planting at 100 lb per acre (F); (S) and (F).

Nomersan, 10% Thiram dust applied: to seed at 3 oz. per cwt (S);
to furrows before planting at 100 lb per acre (F); (S) and (F).

Griseofulvin applied: to seed as 50% glycerol paste at 0.6 g. per
tuber (S); to furrows before planting as 0.4% suspension at
100 gallons per acre (F); to foliage as 0.04% suspension at
50 gallons per acre (L).

Basal dressing: 8 tons dung per acre ploughed in; $10\frac{1}{2}$ cwt compound
granular fertilizer (9% N, 9% P_2O_5 , 15% K_2O) per acre before ridging.

Cultivations, etc.: Dung applied, ploughed in: Sept 28, 1955. Basal
fertilizer applied on flat: Apr 4, 1956. Ridged: Apr 16. Potatoes
hand planted: Apr 17. Earthed up: June 29. Sprayed with copper
fungicide, 5 lb in 40 gallons per acre: July 23. Sprayed again at
3 lb in 30 gallons per acre: Aug 15. Sprayed with sulphuric acid,
20% BOV, 80 gallons per acre: Sept 12. Lifted: Oct 9. Variety:
Majestic. Previous crop: Wheat.

Standard error per plot:

Total tubers: 1.12 tons per acre or 7.5% (30 d.f.)

Note: Emergence counts and records of incidence of root browning and
skin spot (Oospora pustulans), black scurf and stem canker
(Corticium solani) and common scab (Actinomyces scabies) were made.

56/Ce/5.2

Summary of Results

Method of Application	Fungicide					Mean
	None	P	C	N	A	
Total tubers: tons per acre						
			(± 0.649)			(± 0.324)
S		13.67	14.73	15.31	12.95	14.16
F		14.31	16.77	14.97	16.07	15.53
SF		14.69	14.97	14.50		14.72
L					15.48	15.48
Mean (± 0.375)	15.11	14.22	15.49	14.92	14.83	14.92

(1) ± 0.375 (2) ± 0.649 Percentage ware ($1\frac{1}{2}$ " riddle)

S	88.1	84.1	91.4	80.5	86.0
F	85.3	87.2	86.2	88.0	86.7
SF	90.1	85.9	86.5		87.5
L				87.5	87.5
Mean	86.2	87.9	85.7	88.0	85.3

Fungicide

P = PCNB (Parachloronitro benzene) 20% dust.

C = Calomel 4% dust.

N = Nomersan 10% thiram dust.

A = Griseofulvin.

Methods of application

S = Seed treated.

F = Applied to furrows before planting.

L = Applied as spray to foliage.