

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1958

[Full Table of Content](#)



### 58/W/CB/7 Barley - N and Residual Dung N P and K

#### Rothamsted Research

Rothamsted Research (1959) *58/W/CB/7 Barley - N and Residual Dung N P and K* ; Yields Of The Field Experiments 1958, pp 101 - 102 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/Cb/7.1

BARLEY

Residual effects of dung, N, P and K applied to potatoes 1957 and direct effect of N - Great Hill Woburn 1958.

Design: 4 randomized blocks of 16 plots each, certain high order interactions being confounded with block differences.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

To potatoes 1957.

Dung: None; 5; 10; 20 tons per acre ploughed in.  
N: None; 0.9 cwt N per acre as sulphate of ammonia.  
P: None; 0.75 cwt  $P_2O_5$  per acre as superphosphate.  
K: None; 1.5 cwt  $K_2O$  per acre as muriate of potash.

To barley 1958.

N: None; 0.4 cwt N per acre as 'Nitro-Chalk'.

Basal dressing:  $2\frac{1}{2}$  cwt compound fertilizer (16%  $P_2O_5$ , 16%  $K_2O$ ) per acre.

Cultivations, etc.: Ploughed: Dec 10, 1957. 'Nitro-Chalk' and basal fertilizer applied, seed drilled at 3 bushels per acre: Apr 9, 1958. Sprayed with MCPA at 5 pints in 40 gallons per acre: May 20. Combine harvested: Aug 27. Variety: Herta. Previous crop: Potatoes.

Standard error per plot.

Grain (at 85% dry matter): 3.25 cwt per acre or 14.7% (34 d.f.)\*

\*1 missing value.

Summary of Results

Grain (at 85% dry matter): cwt per acre

		Dung to potatoes 1957: tons per acre				Mean
		None	5	10	20	
Mean	(±0.81)	22.0	21.1	22.7	22.8	22.1
<u>Applied to potatoes 1957</u>						
N: cwt per acre						
None	(±1.15)	21.1	20.1	21.8	20.7	20.9
0.9		22.9	22.1	23.6	25.0	23.4
Difference	(±1.62)	+1.8	+2.0	+1.8	+4.3	+2.5 (±0.81)
P <sub>2</sub> O <sub>5</sub> : cwt per acre						
None	(±1.15)	22.9	20.3	21.3	22.9	21.9
0.75		21.0	21.9	24.0	22.7	22.4
Difference	(±1.62)	-1.9	+1.6	+2.7	-0.2	+0.5 (±0.81)
K <sub>2</sub> O: cwt per acre						
None	(±1.15)	22.9	21.2	23.3	23.6	22.7
1.5		21.1	21.0	22.1	22.1	21.5
Difference	(±1.62)	-1.8	-0.2	-1.2	-1.5	-1.2 (±0.81)
<u>Applied to barley 1958</u>						
N: cwt per acre						
None	(±1.15)	16.5	15.1	18.3	16.6	16.6
0.4		27.5	27.1	27.0	29.0	27.7
Difference	(±1.62)	+11.0	+12.0	+8.7	+12.4	+11.1 (±0.81)

Response to	Responses to treatments cwt per acre							
	Applied to potatoes 1957						Applied to barley 1958	
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N	P <sub>2</sub> O <sub>5</sub>
	None	0.9	None	0.75	None	1.5	None	0.4
<u>Applied to potatoes 1957</u>	(±1.15)							
N	-	-	+2.2	+2.8	+2.3	+2.7	+3.3	+1.7
P <sub>2</sub> O <sub>5</sub>	+0.2	+0.8	-	-	+0.4	+0.6	+1.2	-0.2
K <sub>2</sub> O	-1.4	-1.0	-1.3	-1.1	-	-	-1.5	-0.9
<u>Applied to barley 1958</u>								
N	+11.8	+10.2	+11.7	+10.3	+10.7	+11.3	-	-

Mean dry matter % as harvested: 80.7