


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CE/2 Beans - Time of Sowing, Spraying P and K

Rothamsted Research

Rothamsted Research (1959) *58/R/CE/2 Beans - Time of Sowing, Spraying P and K* ; Yields Of The Field Experiments 1958, pp 107 - 108 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

BEANS

Time of sowing, spraying, P and K - Little Knott I 1958.

Design: 3 blocks of 4 whole plots each, plots being split into 3 for P and K with spraying on pairs of whole plots, time of sowing on whole plots, and PK partially confounded.

Area of each sub plot: 0.0337 acres. Area harvested: 0.0105 acres.

Treatments. All combinations of:-

Time of sowing: Autumn; spring.

Spray: None; demeton-methyl (new formulation) at 6 fluid oz of 50% active ingredient in 60 gallons per acre.

Phosphate: None; 0.5; 1.0 cwt P_2O_5 per acre as superphosphate.

Potash: None; 1.0; 2.0 cwt K_2O per acre as muriate of potash.

Basal dressing: None.

Cultivations, etc.: Floughed: Sept 12, 1957. Fertilizers applied for autumn beans, seed drilled at 275 lb per acre: Oct 8. Fertilizers applied for spring beans: Feb 22, 1958. Seed drilled at 200 lb per acre: Mar 7. Appropriate plots sprayed with demeton-methyl: June 18. Combine harvested: Sept 25. Variety: Winter beans - S.Q.Giant, spring beans - Albyn Tick. Previous crop: Spring wheat.

Note. A similar experiment at Woburn was abandoned due to bird damage.

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 1.24 cwt per acre or 9.5% (4 d.f.)

Sub plot: 1.51 cwt per acre or 11.5% (12 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Sown			P ₂ O ₅ : cwt per acre			K ₂ O: cwt per acre			Mean
	Autumn	Spring	Diff.	None	0.5	1.0	None	1.0	2.0	
				(±0.62)*			(±0.62)*			
<u>Spray</u>										
None	12.8	14.4	+1.6	14.8	14.1	11.9	14.3	14.0	12.6	13.6
Demeton-methyl	10.6	14.5	+3.9	13.6	11.6	12.5	13.5	12.8	11.3	12.6

Diff.	-2.2	+0.1	+2.3	-1.2	-2.5	+0.6	-0.8	-1.2	-1.3	-1.0
			(±1.43)	(±0.87)**			(±0.87)**			
				(1) & (2)			(1) & (2)			
<u>Sown</u>										
Autumn				12.9	11.0	11.2	12.3	11.5	11.3	11.7
Spring				15.4	14.7	13.2	15.5	15.3	12.6	14.5

Mean				14.2	12.8	12.2	13.9	13.4	12.0	13.1
(±0.44)										
Diff.				+2.5	+3.7	+2.0	+3.2	+3.8	+1.3	+2.8
(±1.01)										(±0.72)

Mean dry matter % as harvested: Autumn sown, 70.7; spring sown, 72.6.

(1) ±0.71 for use in diagonal comparison only.

(2) ±0.62 for use in horizontal and interaction comparisons only.

* For use in horizontal comparisons only.

** For use only in testing the difference of two differences.