


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CG/2 Grass - N and K

Rothamsted Research

Rothamsted Research (1959) *58/R/CG/2 Grass - N and K* ; Yields Of The Field Experiments 1958, pp 115 - 115 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

GRASS

Levels of N and K - Harwoods Piece 1958.

Design: 4 randomized blocks of 12 plots each.

Area of each plot: 0.0087 acres. Area harvested: 1st cut - 0.0050 acres, 2nd and 3rd cuts - 0.0035 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre as 'Nitro-Chalk'.

Potash: None; 0.3; 0.6 cwt K₂O per acre as muriate of potash.

All treatments in the presence of 0.6 cwt P₂O₅ per acre as superphosphate.

In addition 2 plots per block, receiving 0.9 N and 0.6 K also received phosphate at either None or 1.2 cwt P₂O₅ per acre as superphosphate.

Basal dressing: None.

Cultivations, etc.: Ploughed: Oct 28 - Nov 22, 1957. Fertilizers applied in seedbed: Apr 23, 1958. Seeds sown: Apr 24. Sprayed with MCPA at 4 pints in 40 gallons per acre: May 27. 2nd and 3rd dressings of N and K applied: July 16 and Sept 9. Cut 3 times: July 9, Sept 9 and Oct 27. Variety: S22 Italian Ryegrass. Previous crop: Spring wheat.

Standard error per plot. Dry matter:

1st cut: 2.03 cwt per acre or 7.7% (33 d.f.)
 2nd cut: 1.91 cwt per acre or 6.7% (33 d.f.)
 3rd cut: 0.83 cwt per acre or 5.9% (33 d.f.)
 Total of 3 cuts: 3.06 cwt per acre or 4.5% (33 d.f.)

Summary of Results

Dry matter: cwt per acre

cwt per acre	Dry matter: cwt per acre												
N	0.0	0.3	0.3	0.3	0.6	0.6	0.6	0.9	0.9	0.9	0.9	0.9	
P ₂ O ₅	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.0	1.2	
K ₂ O	0.0	0.0	0.3	0.6	0.0	0.3	0.6	0.0	0.3	0.6	0.6	0.6	Mean
1st cut (±1.01)	15.2	21.6	22.3	19.8	27.3	27.6	27.5	31.6	30.3	30.4	30.5	31.0	26.3
2nd cut (±0.95)	15.6	25.1	25.2	26.2	29.8	30.2	30.1	32.0	33.8	32.1	30.8	32.1	28.6
3rd cut (±0.41)	4.0	12.3	12.7	11.9	14.6	15.5	15.0	17.1	15.5	16.0	15.6	17.4	13.9
Total of 3 cuts (±1.53)	34.8	59.0	60.2	57.9	71.6	73.3	72.6	80.6	79.5	78.4	76.8	80.5	68.8

Mean dry matter % as cut:

1st cut: 15.4 3rd cut: 12.6
 2nd cut: 12.4 Total of 3 cuts: 13.4