


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1958

[Full Table of Content](#)



---

### 58/R/BA/2 and 58/W/BA/2 Six-course Rotation - Cereal Strips

#### Erratum

Erratum (1959) *58/R/BA/2 and 58/W/BA/2 Six-course Rotation - Cereal Strips* ; Yields Of The Field Experiments 1958, pp 26 - 29 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/Ba/2.1

## SIX COURSE ROTATION EXPERIMENT

The 29th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1958.

For history, treatments, etc., see "Details of the Classical and Long Term Experiments" 1956.

Area of each plot: Rothamsted, 0.0250 acres; Woburn, 0.0266 acres.

Cultivations, etc.:

### Rothamsted

Sugar beet.

Ploughed twice: Aug 22 and Nov 2, 1957. Fertilizers applied, seed drilled at 19 lb per acre: Apr 14, 1958. Singled: June 4. Sprayed with demeton methyl, 12 oz in 40 gallons per acre: July 7. Lifted: Nov 24. Variety: Klein E.

Barley.

Sugar beet tops spread: Nov 21, 1957. Ploughed: Nov 22. Ground chalk applied at 23 cwt per acre: Nov 23. Fertilizers applied: Mar 21, 1958. Seed drilled at  $2\frac{3}{4}$  bushels per acre: Mar 22. Clover seed undersown: Apr 25. Harvested: Aug 27. Variety: Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: May 20, 1957. Autumn fertilizers applied: Sept 23. Sulphate of ammonia applied: Apr 8, 1958. Cut: July 8. Variety: S123 Late Flowering Red.

Wheat<sup>‡</sup>

Ploughed twice: July 7 and Oct 9, 1957. Autumn fertilizers applied, seed drilled at  $2\frac{3}{4}$  bu. per acre: Oct 14. Sulphate of ammonia applied: Apr 9, 1958. Sprayed with CMFP, 6 pints in 40 gallons per acre: Apr 30. Harvested: Sept 3 - 11. Variety: Yeoman.

Potatoes.

Ploughed twice: Aug 22 and Oct 9, 1957. Ridged, fertilizers applied, potatoes planted: Apr 19, 1958. Earthed up: July 8. Sprayed 4 times with copper fungicide, 5 lb in 40 gallons per acre: July 12 and at 3 lb and 1 pint of spreader in 40 gallons per acre: Aug 1, 6 and 16. Sprayed with sulphuric acid, 20% BOV at 100 gallons per acre: Sept 9. Lifted: Sept 25. Variety: Majestic.

Rye.

Ploughed: Oct 9, 1957. Ground chalk applied at 23 cwt per acre: Oct 14. Autumn fertilizers applied, seed drilled at 3 bushels per acre: Oct 15. Sulphate of ammonia applied: Apr 9, 1958. Sprayed with CMFP, 6 pints in 40 gallons per acre: Apr 30. Harvested: Aug 26. Variety: King II.

<sup>‡</sup> Owing to severe lodging, yields from this crop were estimated from a 6 ft. wide sample cut from each plot (Area 0.0054 acres).

58/Ba/2.2

Woburn

Sugar beet.

Ploughed twice: Aug 29 and Nov 30, 1957. Fertilizers applied, seed drilled at 10 lb per acre: Apr 17, 1958. Sprayed with miscible DDT at 3 pints in 40 gallons per acre: May 3. Singled: June 6. Sprayed with demeton methyl, 12 oz in 40 gallons per acre: June 26. Lifted: Oct 15. Variety: Klein E.

Barley.

Ploughed: Nov 29, 1957. Fertilizers applied: Mar 19, 1958. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Mar 20. Harvested: Aug 25. Variety: Herta.

Clover.

Ploughed: Aug 9 and 17, 1957. PK fertilizers applied: Aug 30. Seed broadcast at 40 lb per acre: Sept 5. Ploughed: Mar 4, 1958. Nitrogen fertilizers applied: Mar 19. Resown: Mar 24. Cut: July 14. Variety: Crimson Clover.

Wheat.

Ploughed twice: July 8 and Aug 29, 1957. Autumn fertilizers applied: Oct 17. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Oct 26. Sulphate of ammonia applied: Apr 21, 1958. Sprayed with CMPP, 6 pints in 40 gallons per acre: May 3. Harvested: Aug 26. Variety: Yeoman.

Potatoes.

Ploughed twice: Aug 29, 1957 and Jan 14, 1958. Fertilizers applied, and potatoes hand planted: Apr 21. Earthed up: June 18. Sprayed with copper fungicide, 5 lb in 80 gallons per acre: July 14. Twice sprayed with copper fungicide, 5 lb in 40 gallons per acre: July 29 and Aug 16. Sprayed with arsenious compound, 1 gallon in 40 gallons per acre: Sept 12. Lifted: Oct 7. Variety: Majestic.

Rye.

Ploughed: Oct 5, 1957. Ground chalk applied at 20 cwt per acre: Oct 10. Fertilizers applied: Oct 17. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Oct 26. Sprayed with CMPP, 6 pints in 40 gallons per acre: May 3, 1958. Sulphate of ammonia applied: Apr 21. Harvested: Aug 26. Variety: King II.

58/Ba/2.3

Summary of Results

Mean yields per acre and responses in yield per cwt of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O

	Rothamsted	Woburn	Rothamsted	Woburn
Sugar Beet, roots (washed): tons per acre			Barley, grain: cwt per acre	
Mean	14.24	8.77	24.5 <sup>‡</sup>	26.1 <sup>‡</sup>
Response to: N	+6.52	+4.22	+9.4	+4.0
P	+1.25	+7.69	-3.5	-7.9
K	-1.28	+1.02	+1.2	+3.3
Mean dry matter % as harvested:			78.9	81.1
Sugar Beet sugar percentage			Barley, straw: cwt per acre	
Mean	17.4	16.6	29.6 <sup>‡</sup>	32.3 <sup>‡</sup>
Response to: N	-0.4	+0.2	+6.6	+14.9
P	-1.0	+0.1	+11.5	+1.7
K	+0.2	+0.4	-0.6	+0.6
Mean dry matter % as harvested:			72.2	83.2
Sugar Beet, total sugar: cwt per acre			Clover, hay, dry matter: cwt per acre	
Mean	49.6	29.4	47.7	14.9
Response to: N	+21.3	+14.1	-7.0	+10.7
P	+1.5	+25.8	-5.9	+13.3
K	-3.6	+4.1	+1.2	-6.2
Mean dry matter % as cut:			60.2	17.9
Sugar Beet, tops: tons per acre			Wheat, grain: cwt per acre	
Mean	9.17	7.10	18.7 <sup>‡</sup>	20.6 <sup>‡</sup>
Response to: N	+3.27	+4.42	-6.7	+19.3
P	-0.24	+5.93	-3.4	+1.8
K	+0.80	-1.40	+2.6	+0.6
Mean dry matter % as harvested:			77.2	80.1
Sugar Beet, plant number: thousands per acre			Wheat, straw: cwt per acre	
Mean	31.1	‡‡‡	46.9 <sup>‡</sup>	28.4 <sup>‡</sup>
Response to: N	+0.4		+14.1	+31.0
P	-0.3		-7.9	+3.4
K	-1.0		+8.0	+0.1
Mean dry matter % as harvested:			75.5	82.4

<sup>‡</sup> (At 85% dry matter).      <sup>‡‡‡</sup> Not recorded.

58/Ba/2.4

Mean yields per acre and responses in yield per cwt of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O

	Rothamsted	Woburn	Rothamsted	Woburn
	Potatoes, total tubers tons per acre		Rye, grain: cwt per acre	
Mean	7.75	6.48	23.2 <sup>‡</sup>	25.2 <sup>‡</sup>
Response to: N	+8.41	+4.04	+11.6	+14.4
P	+1.85	-2.91	-3.9	+3.3
K	+0.32	+3.23	+1.4	-0.6
Mean dry matter % as harvested:			81.2	80.4
	Potatoes, percentage ware		Rye, straw: cwt per acre	
	(1)	(2)		
Mean	87.2	75.0	41.5 <sup>‡</sup>	34.1 <sup>‡</sup>
Response to: N	+6.0	+0.2	+31.0	+19.8
P	-7.7	+10.0	-11.3	+6.4
K	+0.9	+5.8	+4.8	-3.3
Mean dry matter % as harvested:			83.4	82.3

<sup>‡</sup> (At 85% dry matter)

Riddle: (1) 1½"; (2) 1⅝"