


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CA/1 Winter Wheat - Seed Rates, Sowing Dates and N

Rothamsted Research

Rothamsted Research (1959) *58/R/CA/1 Winter Wheat - Seed Rates, Sowing Dates and N* ; Yields Of The Field Experiments 1958, pp 76 - 77 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/Oa/1.1

WINTER WHEAT

Seed rates, sowing dates and levels of nitrogen - Great Field I 1958.

Design: 3 randomized blocks of 9 plots, each plot being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0148 acres. Area harvested: 0.0096 acres.

Treatments. All combinations of:-

Seed rates: 2; 3; 4 bushels per acre.

Sowing dates: Sept 17; Oct 7; Nov 8, 1957.

Nitrogen: 0.6; 1.2 cwt N per acre applied as 'Nitro-Chalk' in two equal parts in early February and late April.

Basal dressing: 3 cwt compound fertilizer (10% P_2O_5 , 20% K_2O) per acre broadcast in seed bed, 3 cwt compound fertilizer (5% N, $12\frac{1}{2}\%$ P_2O_5 , $12\frac{1}{2}\%$ K_2O) per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Aug 21, 1957. Compound fertilizer applied by hand: First sowing - Sept 17; second sowing - Oct 7; third sowing - Nov 8. First dressing of N applied: Feb 15, 1958. Sprayed with CMPP at 6 pints in 40 gallons per acre: Apr 17. Second dressing of N applied: Apr 24. Combine harvested: Sept 10. Variety: Cappelle. Previous crop: Potatoes.

Standard errors per plot. Grain (at 85% dry matter):

Whole plot: 2.36 cwt per acre or 7.7% (16 d.f.)

Sub plot: 2.25 cwt per acre or 7.3% (18 d.f.)

Note. Estimates of % area lodged and counts of plant shoot and ear number were made. All plots were completely lodged at harvest.

58/Ca/1.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Seed rate: bushels per acre	Date of sowing			N: cwt per acre		Diff.	Mean
	Sept 17th	Oct 7th	Nov 8th	0.4	0.8		
		(±0.96)		(±0.95)*		(±1.06)	(±0.56)
2	33.2	31.1	34.3	35.0	30.7	-4.3	32.8
3	29.3	27.2	33.7	30.8	29.3	-1.5	30.1
4	31.2	29.2	26.9	29.8	28.4	-1.4	29.1
			Date of sowing	(±0.95)*		(±1.06)	(±0.56)
			Sept 17th	31.3	31.1	-0.2	31.2
			Oct 7th	30.1	28.1	-2.0	29.1
			Nov 8th	34.1	29.2	-4.9	31.6
			Mean	31.8	29.5	-2.3	30.7
						(±0.61)	

*For use in vertical and diagonal comparisons.

Mean dry matter % as harvested: 74.1