


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CA/7 and 58/W/CA/7 Spring Wheat - Varieties and N

Rothamsted Research

Rothamsted Research (1959) *58/R/CA/7 and 58/W/CA/7 Spring Wheat - Varieties and N* ; Yields Of The Field Experiments 1958, pp 86 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/Ca/7.1

SPRING WHEAT

Varieties and levels of nitrogen - Rothamsted (R) Great Knott III and Woburn (W) Stackyard 1958.

Design (each field): 3 randomized blocks of 7 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot (acres):	Area harvested (acres):
Great Knott III (R): 0.0154	0.0103
Stackyard (W): 0.0193	0.0129

Treatments: All combinations of:-

Whole plots. Varieties: Atle (1); Atson (2); Koga II (3); Miana (4); Peko (5); Progress (6); Svenno (7).

Sub plots: Nitrogen, applied as sulphate of ammonia.

Great Knott III (R): None; 0.4 cwt N per acre
Stackyard (W): 0.4; 0.8 cwt N per acre.

Basal dressing:

Great Knott III (R): 3.3 cwt compound fertilizer (12% N, 9% P₂O₅, 9% K₂O) per acre combine drilled with seed.

Stackyard (W): 1 cwt compound fertilizer (16% P₂O₅, 16% K₂O) per acre combine drilled with seed.

Cultivations, etc.:

Great Knott III (R). Ploughed: Oct 28, 1957. Seed combine drilled at 3¼ bushels per acre: Apr 4, 1958. Nitrogen applied: Apr 8. Sprayed with MCPA at 4 pints in 40 gallons per acre: Apr 20. Combine harvested: Sept 12. Previous crop: Spring wheat.

Stackyard (W). Ploughed: Feb 13, 1958. Nitrogen applied: Mar 24. Seed combine drilled at 3¼ bushels per acre: Apr 2. Sprayed with MCPA at 4 pints in 40 gallons per acre: May 27. Combine harvested: Oct 9. Previous crop: Potatoes.

Standard errors per plot, Grain (at 85% dry matter):

Great Knott III (R):

Whole plot: 1.53 cwt per acre or 6.2% (12 d.f.)

Sub plot: 1.98 cwt per acre or 8.0% (14 d.f.)

Stackyard (W):

Whole plot: 1.61 cwt per acre or 6.2% (12 d.f.)

Sub plot: 1.85 cwt per acre or 7.1% (14 d.f.)

58/Ca/7.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre (including basal)	Variety							Mean
	1	2	3	4	5	6	7	
	Great Knott III (R)							
	(±1.20)*							
0.4	24.6	24.1	23.4	15.4	23.2	23.6	23.3	22.5
0.8	24.6	26.0	26.8	25.1	26.8	25.9	31.0	26.6
Mean (±0.88)	24.6	25.0	25.1	20.2	25.0	24.8	27.1	24.5
Difference (±1.62)	0.0	1.9	3.4	9.7	3.6	2.3	7.7	4.1 (±0.61)

Mean dry matter % as harvested: 80.6

N: cwt per acre	Stackyard (W)							Mean
	(±1.20)*							
0.4	23.4	22.9	26.2	21.3	26.2	25.1	23.7	24.1
0.8	28.1	26.3	28.9	24.9	29.3	30.5	25.9	27.7
Mean (±0.93)	25.8	24.6	27.5	23.1	27.7	27.8	24.8	25.8
Difference (±1.51)	4.7	3.4	2.7	3.6	3.1	5.4	2.2	3.6 (±0.57)

Mean dry matter % as harvested: 73.6

*for use in comparisons other than vertical.