

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/A/4 Barnfield - Mangolds and Sugar Beet

Rothamsted Research

Rothamsted Research (1959) *58/R/A/4 Barnfield - Mangolds and Sugar Beet* ; Yields Of The Field Experiments 1958, pp 9 - 10 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

58/A/4.1

MANGOLDS AND SUGAR BEET - BARNFIELD

The 83rd and 13th years

For history, treatments etc., see "Details of the Classical and Long Term Experiments" 1956.

Cultivations, etc.: Dung applied, ploughed: Nov 25, 1957.
 Fertilizers applied: May 5, 1958. Sugar beet drilled at 18 lb per acre, mangolds drilled at 8 lb per acre: May 8. Singled: June 11 to July 5. Sprayed with demeton methyl at 12 oz in 40 gallons per acre: July 7. Top dressings applied: July 14. Lifted: Oct 11 - Nov 21. Varieties: Mangolds - Yellow Globe, sugar beet - Klein E.

Summary of Results

Strip	Cross Dressing				
	O	N	A	AC	C
Mangolds, roots: tons per acre					
1	12.74	31.56	28.54	24.15	17.27
2	17.06	36.47	30.34	30.34	25.70
4	6.38	(a) 26.80* (b) 27.47*	22.13	28.63	20.34
5	4.93	20.44	16.17	18.64	14.51
6	4.82	19.73	19.68	26.63	20.84
7	5.84	21.52	20.44	26.87	21.06
8	1.93	16.34	13.95	19.70	15.27
9	19.80				
Mangolds, leaves: tons per acre					
1	2.42	5.69	5.59	6.23	4.01
2	2.69	6.62	5.54	8.04	5.67
4	1.32	(a) 5.52* (b) 5.01*	3.64	6.77	4.27
5	0.98	4.03	3.35	3.83	3.15
6	1.00	3.76	3.15	4.67	4.23
7	1.34	4.79	3.96	5.64	4.93
8	0.81	3.93	3.76	4.20	4.52
9	3.81				
Mangolds, plant number: thousands per acre					
1	18.9	19.7	19.1	18.0	17.7
2	23.6	20.8	20.5	19.5	20.7
4	21.8	(a) 21.8* (b) 21.1*	20.3	19.7	20.2
5	21.6	21.3	21.9	20.7	21.5
6	20.5	21.5	20.7	20.2	21.0
7	20.9	21.8	21.0	20.4	21.3
8	19.0	20.5	21.5	21.4	21.6
9	24.0				

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

58/A/4.2

Strip	Cross Dressing				
	O	N	A	AC	C
Sugar beet, roots (washed): tons per acre					
1	9.04	13.41	13.47	14.26	10.84
2	6.85	12.69*	12.62	13.65	12.47
4	2.08	(b) 11.37*	10.43	14.31	12.75
5	2.77	8.76	9.29	10.76	9.84
6	1.89	9.95	10.32	13.65	11.83
7	2.43	10.30	10.60	12.64	10.81
8	1.89	7.41	7.56	11.24	9.22
9	8.39				
Sugar beet, tops: tons per acre					
1	6.64	13.14	16.61	17.00	15.29
2	6.20	14.85*	14.36	18.76	13.48
4	1.81	(b) 12.02*	6.50	14.70	11.92
5	1.61	10.11	7.82	15.88	10.11
6	1.66	10.60	6.55	15.09	10.70
7	1.91	13.24	9.62	17.05	11.14
8	1.86	9.67	7.57	15.24	11.87
9	7.43				
Sugar beet, plant number: thousands per acre					
1	23.7	22.7	23.4	22.8	22.9
2	24.3	22.8*	23.5	24.4	24.3
4	22.3	(b) 24.0*	24.5	23.6	24.9
5	23.0	23.7	23.5	22.2	23.9
6	22.7	24.3	24.9	24.8	25.2
7	22.3	24.3	23.9	23.3	23.3
8	22.4	23.0	24.4	25.0	25.0
9	22.6				
Sugar beet, sugar percentage					
1	17.9	17.2	17.0	17.0	16.8
2	17.7	17.0*	16.6	16.4	17.0
4	18.0	(b) 17.4*	18.4	17.4	17.6
5	17.7	16.9	17.8	16.5	17.1
6	17.4	17.2	18.2	17.0	17.4
7	17.8	17.6	18.3	16.9	17.5
8	17.4	17.3	18.0	17.2	17.3
9	17.9				

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.