


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CB/2 and 58/W/CB/2 Barley - N and Residual Dung N P K and Salt

Rothamsted Research

Rothamsted Research (1959) *58/R/CB/2 and 58/W/CB/2 Barley - N and Residual Dung N P K and Salt* ; Yields Of The Field Experiments 1958, pp 90 - 92 - DOI:

<https://doi.org/10.23637/ERADOC-1-181>

58/Cb/2.1

BARLEY

Residual effects of dung, N, P, K and salt to sugar beet 1957 and direct effect of N - Rothamsted (R) West Barnfield II and Woburn (W) Great Hill 1958.

Design: Half replicate of 4×2^5 arranged in 4 blocks of 16 plots each, the identity being $d(20 - 10 - 5 + 0)npksN = 1$, with certain high order interactions confounded with block differences.

Area of each plot (each field): 0.0167 acres. Area harvested: 0.0141 acres.

Treatments: All combinations of:-

Applied to sugar beet 1957

Dung: None; 5; 10; 20 tons per acre ploughed in.
N: None; 0.9 cwt N per acre as sulphate of ammonia.
P: None; 0.75 cwt P_2O_5 per acre as superphosphate.
K: None; 1.5 cwt K_2O per acre as muriate of potash.
Salt: None; 5 cwt agricultural salt per acre.

Applied to barley 1958

N: None; 0.4 cwt N per acre as 'Nitro-Chalk'.

Basal dressing (each field): $2\frac{1}{2}$ cwt compound fertilizer (16% P_2O_5 , 16% K_2O) per acre combine drilled with seed.

Cultivations, etc.:

West Barnfield II (R). Ploughed: Jan 1. Seed combine drilled with basal fertilizers at 2 bushels per acre, nitrogen fertilizer applied: Mar 25. Sprayed with MCPB at 5 pints in 40 gallons per acre: May 28. Combine harvested: Aug 27. Variety: Proctor. Previous crop: Sugar beet.

Great Hill (W). Ploughed: Jan 27. Nitrogen fertilizer applied, seed combine drilled at 3 bushels per acre with basal fertilizer: Apr 9. Sprayed with MCPA at 5 pints in 40 gallons per acre: May 20. Combine harvested: Aug 26. Variety: Herta. Previous crop: Sugar beet.

Standard errors per plot, Grain (at 85% dry matter):

West Barnfield II (R): 1.48 cwt per acre or 3.9% (27 d.f.)
Great Hill (W): 1.48 cwt per acre or 5.8% (27 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Dung to sugar beet 1957:				Mean
	None	5	10	20	
West Barnfield II (R)					
Mean (± 0.37)	36.4	37.6	37.8	38.8	37.7
<u>Applied to sugar beet 1957</u>					
N: cwt per acre					
None (± 0.52)	35.4	37.5	37.9	38.5	37.3
0.9 (± 0.52)	37.4	37.8	37.7	39.1	38.0
Difference (± 0.74)	+2.0	+0.3	-0.2	+0.6	+0.7 (± 0.37)
P ₂ O ₅ : cwt per acre					
None (± 0.52)	36.0	37.4	37.1	38.1	37.1
0.75 (± 0.52)	36.8	37.9	38.5	39.5	38.2
Difference (± 0.74)	+0.8	+0.5	+1.4	+1.4	+1.1 (± 0.37)
K ₂ O: cwt per acre					
None (± 0.52)	37.2	37.6	37.9	38.7	37.9
1.5 (± 0.52)	35.7	37.6	37.7	38.9	37.5
Difference (± 0.74)	-1.5	0.0	-0.2	+0.2	-0.4 (± 0.37)
Salt: cwt per acre					
None (± 0.52)	36.6	37.7	38.1	38.8	37.8
5.0 (± 0.52)	36.3	37.5	37.5	38.9	37.5
Difference (± 0.74)	-0.3	-0.2	-0.6	+0.1	-0.3 (± 0.37)
<u>Applied to barley 1958</u>					
N: cwt per acre					
None (± 0.52)	33.7	35.7	36.3	37.3	35.7
0.4 (± 0.52)	39.1	39.5	39.4	40.4	39.6
Difference (± 0.74)	+5.4	+3.8	+3.1	+3.1	+3.9 (± 0.37)

Response to	Responses to treatments								Applied to barley 1958	
	cwt per acre									
	Applied to sugar beet 1957				Applied to sugar beet 1957				N	
	None	N 0.9	P ₂ O ₅ None	P ₂ O ₅ 0.75	K ₂ O None	K ₂ O 1.5	Salt None	Salt 5.0	None	0.4
<u>Applied to sugar beet 1957</u>										
N	-	-	+1.3	+0.1	+1.1	+0.3	+1.1	+0.3	+2.1	-0.7
P ₂ O ₅	+1.7	+0.5	-	-	+1.4	+0.8	+0.6	+1.6	+1.3	+0.9
K ₂ O	0.0	-0.8	-0.1	-0.7	-	-	-0.2	-0.6	-0.3	-0.5
Salt	+0.1	-0.7	-0.8	+0.2	-0.1	-0.5	-	-	-0.5	-0.1
<u>Applied to barley 1958</u>										
N	+5.2	+2.4	+4.0	+3.6	+3.9	+3.7	+3.6	+4.0	-	-
Mean dry matter % as harvested:	80.7									

58/Oh/2.3

Grain (at 85% dry matter): cwt per acre

		Dung to sugar beet 1957: tons per acre				Mean
		None	5	10	20	
Great Hill (W)						
Mean	(±0.37)	25.0	25.6	26.0	25.5	25.5
<u>Applied to sugar beet 1957</u>						
N: cwt per acre						
None	(±0.52)	24.4	25.2	25.9	25.5	25.3
0.9		25.6	25.9	26.2	25.5	25.8
Difference	(±0.74)	+1.2	+0.7	+0.3	0.0	+0.5 (±0.37)
P ₂ O ₅ : cwt per acre						
None	(±0.52)	25.2	25.1	26.1	25.2	25.4
0.75		24.8	26.1	26.0	25.8	25.7
Difference	(±0.74)	-0.4	+1.0	-0.1	+0.6	+0.3 (±0.37)
K ₂ O: cwt per acre						
None	(±0.52)	24.7	25.9	26.1	25.8	25.6
1.5		25.2	25.2	26.0	25.3	25.4
Difference	(±0.74)	+0.5	-0.7	-0.1	-0.5	-0.2 (±0.37)
Salt: cwt per acre						
None	(±0.52)	24.6	26.0	25.7	25.3	25.4
5.0		25.4	25.1	26.4	25.7	25.6
Difference	(±0.74)	+0.8	-0.9	+0.7	+0.4	+0.2 (±0.37)
<u>Applied to barley 1958</u>						
N: cwt per acre						
None	(±0.52)	19.4	19.8	20.4	19.8	19.8
0.4		30.5	31.4	31.7	31.2	31.2
Difference	(±0.74)	+11.1	+11.6	+11.3	+11.4	+11.4 (±0.37)

Responses to treatments
cwt per acre

Response to	Applied to sugar beet 1957								Applied to barley 1958	
	N	P ₂ O ₅		K ₂ O		Salt		N		
	None	0.9	None	0.75	None	1.5	None	5.0	None	0.4

<u>Applied to sugar beet 1957</u>										
(±0.52)										
N	-	-	+0.1	+0.9	-0.3	+1.3	+0.6	+0.4	+0.9	+0.1
P ₂ O ₅	-0.1	+0.7	-	-	+0.6	0.0	+0.9	-0.3	+0.2	+0.4
K ₂ O	-1.0	+0.6	+0.1	-0.5	-	-	-0.2	-0.2	+0.4	-0.8
Salt	+0.3	+0.1	+0.8	-0.4	+0.2	+0.2	-	-	-0.2	+0.6

<u>Applied to barley 1958</u>										
N	+11.8	+11.0	+11.3	+11.5	+12.0	+10.8	+11.0	+11.8	-	-

Mean dry matter % as harvested: 80.0