


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/R/CB/3 and 58/W/CB/3 Barley - Combine Drilling of N

Rothamsted Research

Rothamsted Research (1959) *58/R/CB/3 and 58/W/CB/3 Barley - Combine Drilling of N* ; Yields Of The Field Experiments 1958, pp 93 - 94 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

BARLEY

Combine drilling of nitrogen - Rothamsted (R) Great Knott III and Woburn (W) Lansome Field 1958.

Design (each field): 4 randomized blocks of 7 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.22* (N₁); 0.54 (N₂); 0.72* (N₃) cwt N per acre.

Method of application: Broadcast as sulphate of ammonia; combine drilled as compound fertilizer:

*N₁: 5% N, 12% P₂O₅, 12½% K₂O
N₂: 8% N, 8% P₂O₅, 8% K₂O
*N₃: 12% N, 9% P₂O₅, 9% K₂O

*Note: For N₁ and N₃ on Great Knott III (R), the compound fertilizer was actually applied at the rates of 0.19 and 0.57 cwt N per acre. The broadcast dressings were corrected to these amounts.

Basal dressing per acre (each field): 0.54 cwt P₂O₅ and 0.54 cwt K₂O combine drilled

(a) as compound (16% P₂O₅, 16% K₂O) on the no nitrogen and broadcast nitrogen plots.

(b) as compounds N₁, N₂, N₃ on the plots receiving drilled nitrogen.

Cultivations, etc.:

Great Knott III (R). Ploughed: Oct 28, 1957. Seed combine drilled at 2¾ bushels per acre: Apr 3, 1958. Sulphate of ammonia broadcast: Apr 4. Sprayed with MCPA at 4 pints in 40 gallons per acre: May 20. Combine harvested: Aug 25. Variety: Herta. Previous crop: Spring wheat.

Lansome Field (W). Ploughed: Nov 27, 1957. Seed combine drilled at 2¾ bushels per acre: Apr 3, 1958. Sulphate of ammonia broadcast: Apr 8. Sprayed with MCPA at 5 pints in 40 gallons per acre: May 22. Combine harvested: Aug 27. Variety: Herta. Previous crop: Fallow.

Standard errors per plot, Grain (at 85% dry matter):

Great Knott III (R): 2.25 cwt per acre or 7.3% (18 d.f.)

Lansome Field (W): 1.79 cwt per acre or 5.8% (18 d.f.)

58/Cb/3.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre

None	Broadcast			Combine drilled			Mean
	0.22 ⁺	0.54	0.72 [*]	0.22 ⁺	0.54	0.72 [*]	

Great Knott III, Rothamsted

19.0	25.0	36.1	36.4 (±1.13)	25.8	37.3	37.2	31.0
------	------	------	-----------------	------	------	------	------

Mean dry matter % as harvested: 74.2

Lansome Field, Woburn

20.0	27.7	33.9	33.4 (±0.89)	28.0	36.5	35.0	30.6
------	------	------	-----------------	------	------	------	------

Mean dry matter % as harvested: 84.3

⁺ 0.19 on Great Knott III, Rothamsted.

^{*} 0.57 " " " " " "