


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1958

[Full Table of Content](#)



58/W/CB/6 Barley - K and Residual N Fertilizers

Rothamsted Research

Rothamsted Research (1959) *58/W/CB/6 Barley - K and Residual N Fertilizers* ; Yields Of The Field Experiments 1958, pp 99 - 100 - DOI: <https://doi.org/10.23637/ERADOC-1-181>

BARLEY

Residual effects of nitrogenous fertilizers and direct effect of potash
- Long Hoos I, II and III 1958.

Design: 4 randomized blocks of 16 plots each, plots being split into 2
for the application of potash.

Area of each sub plot: 0.0038 acres. Area harvested: 0.0032 acres.

Treatments: All combinations of:-

Applied to ryegrass. (whole plots):

None (2 plots per block) together with all combinations of:-

Materials and methods of application

Applied in 1 single dressing:

Formalized casein,	12.2% N in 1956
" "	12.2% N in 1957
Casein,	12.6% N in 1956 and 1957

Applied in 4 dressings of one quarter the single rate:

Ammonium sulphate,	21.0% N in 1957
" "	21.0% N in 1956 and 1957
Urea,	43.5% N in 1956 and 1957
Calcium nitrate,	15.5% N in 1956 and 1957

Rates of application

0.75; 1.5 cwt N per acre

Applied to barley (1958). (sub plots):

K: None; 1.0 cwt K₂O per acre as muriate of potash.

Basal dressing: 3 cwt compound fertilizer (9% N, 18% P₂O₅) per acre
combine drilled with seed.

Cultivations, etc.: Ploughed: Nov 28, 1957. Seed combine drilled at
2½ bushels per acre, muriate of potash applied: Mar 22, 1958.

Sprayed with MCPA at 5 pints in 40 gallons per acre: May 27.

Harvested: Aug 11. Variety: Proctor. Previous crop: Ryegrass.

Standard errors per plot. Grain (at 85% dry matter):

Whole plot: 2.10 cwt per acre or 7.8% (46 d.f.)

Sub plot: 1.76 cwt per acre or 6.6% (49 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Level of N in fertilizer: cwt per acre	Fertilizer								Mean
	None	Single dressing			Divided dressing				
		1956 F	1957 F	1957 & C	1957 A	1956 A	and U	1957 N	
Mean (± 0.74)	25.5	25.4	31.9	26.2	25.7	24.7	27.9	26.9	26.8
				(± 1.05)					
0.75	-	24.2	30.9	26.3	25.1	23.8	27.0	23.5	25.8
1.50	-	26.7	33.0	26.2	26.4	25.6	28.9	30.3	28.2
Diff. (± 1.48)		+2.5	+2.1	-0.1	+1.3	+1.8	+1.9	+6.8	+2.4
				(± 0.86)*					(± 0.56)
K ₂ O: cwt per acre									
None	25.9	25.0	31.5	25.3	25.6	23.7	28.0	26.0	26.4
1.0	25.1	25.8	32.4	27.1	25.9	25.7	27.9	27.8	27.2
Diff. (± 0.88)	-0.8	+0.8	+0.9	+1.8	+0.3	+2.0	-0.1	+1.8	+0.8
									(± 0.31)

Straw (at 85% dry matter): cwt per acre

Level of N in fertilizer: cwt per acre	None	F	C	A	U	N	Mean
Mean	21.6	23.4	29.7	23.5	22.8	22.4	25.4
0.75	-	22.2	27.0	25.1	23.2	21.6	23.6
1.50	-	24.5	32.4	22.0	22.5	23.3	27.2
Diff.		+2.3	+5.4	-3.1	-0.7	+1.7	+3.6
K ₂ O: cwt per acre							
None	22.2	22.9	29.2	22.0	21.8	21.8	25.3
1.0	21.0	23.8	30.2	25.0	23.9	23.1	25.5
Diff.	-1.2	+0.9	+1.0	+3.0	+2.1	+1.3	+0.2
							+2.4

Treatments

- F = Formalized casein 12.2% N
- C = Casein 12.6% N
- A = Ammonium sulphate 21.0% N
- U = Urea 43.5% N
- N = Calcium nitrate 15.5% N

Mean dry matter % as harvested:
Grain, 77.3
Straw, 67.0

*For use in horizontal and diagonal comparisons only.