


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/R/CG/1 Grass - N Fertilizers

Rothamsted Research

Rothamsted Research (1960) *59/R/CG/1 Grass - N Fertilizers* ; Yields Of The Field Experiments 1959, pp 100 - 102 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Cg/1.2

Summary of Results

Rate of application of N: cwt per acre	Dry matter: cwt per acre		Fertilizer		Nitro-Chalk' applied		Mean
	Ureaformaldehyde applied		As single dressing & spring 1959		As divided dressing 1958 & 1959		
	1958	1959	1958 & 1959	1959	1958	1959	
None							
1.0	8.6	13.1	16.9	30.2	9.6	18.6	17.1 (± 0.33)
2.0	9.7	16.7	24.5	33.7	15.3	26.7	22.7
Mean (± 0.62)	9.2	14.9	20.7	32.0	12.4	22.6	18.1
Diff. (± 1.23)	+1.1	+3.6	+7.6	+3.5	+5.7	+8.1	+5.6 (± 0.47)
			<u>1st cut</u>	(± 0.87)			
			<u>2nd cut</u>	(± 1.42)			
1.0	13.1	17.2	14.3	19.5	12.8	26.4	19.1 (± 0.53)
2.0	13.5	22.5	23.1	34.6	13.1	36.7	25.7
Mean (± 1.00)	13.3	19.8	18.7	27.0	12.9	31.5	20.9
Diff. (± 2.00)	+0.4	+5.3	+8.8	+15.1	+0.3	+10.3	+6.6 (± 0.76)

Mean dry matter % as cut:
 1st cut: 17.3
 2nd cut: 32.5

59/Cg/2.1

GRASS

Levels of N and K - Harwoods Piece 1959 - the 2nd year.

Design: 4 randomized blocks of 12 plots each.

Area of each plot: 0.0087 acres. Area harvested: 0.0035 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre as 'Nitro-Chalk'.

Potash: None; 0.3; 0.6 cwt K_2O per acre as muriate of potash.

All treatments in the presence of 0.6 cwt P_2O_5 per acre as superphosphate.

In addition 2 plots per block, receiving 0.9 N and 0.6 K, also received phosphate at either None or 1.2 cwt P_2O_5 per acre as superphosphate.

Note (1): N and K dressings are applied for each cut. All P dressings are applied once annually.

Basal dressing: None.

Cultivations, etc.: 1st dressing of fertilizers applied: Mar 10.

2nd and 3rd dressings of N and K applied: Apr 28 and June 22.

Cut 3 times: Apr 27, June 17 and Aug 18. Variety: S22 Italian Ryegrass.

Standard errors per plot. Dry matter:

1st cut: 1.48 cwt per acre or 5.8% (33 d.f.)

2nd cut: 2.38 cwt per acre or 6.2% (33 d.f.)

3rd cut: 2.48 cwt per acre or 12.3% (33 d.f.)

Total of 3 cuts: 5.26 cwt per acre or 6.3% (33 d.f.)

Note (2): For details of the previous year's results see 'Results of the Field Experiments' 58/Cg/2 to which should be added note (1) and the asterisks and footnote in the summary.