


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1959

[Full Table of Content](#)



59/W/BD/1 Green Manuring

Rothamsted Research

Rothamsted Research (1960) *59/W/BD/1 Green Manuring ; Yields Of The Field Experiments 1959*, pp 47 - 49 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Bd/1.2

Summary of ResultsEarly potatoes, total tubers: tons per acre

	Straw: tons per acre		N: cwt per acre (including basal)		Dung to cabbages 1953: tons per acre		Mean
	None	1½	0.6	1.2	None	10	

Excluding plots fallow under old scheme

Undersown green manures for potatoes	(±0.185)		(±0.185)		(±0.185)		(±0.131)
None	4.12	4.05	4.01	4.16	4.02	4.15	4.08
	(±0.262)		(±0.262)		(±0.262)		(±0.185)
Trefoil	4.21	4.40	4.37	4.24	4.22	4.38	4.30
Ryegrass	4.63	4.86	4.88	4.61	4.28	5.22	4.74
Straw: tons per acre			(±0.185)		(±0.185)		(±0.131)
None			4.24	4.29	4.09	4.44	4.27
1½			4.39	4.29	4.18	4.50	4.34
N: cwt per acre (including basal)					4.10	4.54	4.32
0.6					4.17	4.41	4.29
1.2							
Mean (±0.131)					4.13	4.47	4.30

Plots fallow under old scheme

Straw: tons per acre		(±0.370)		(±0.370)		(±0.262)
None		3.52	3.56	3.42	3.66	3.54
1½		3.96	4.07	3.84	4.19	4.02
N: cwt per acre (including basal)						
0.6				3.58	3.90	3.74
1.2				3.68	3.95	3.82
Mean (±0.262)				3.63	3.92	3.78

Old scheme	Undersown green manures for potatoes				Mean
	None Fallow	None Excluding fallow	Trefoil	Ryegrass	
	3.78 (±0.185)	4.08 (±0.131)	4.30 (±0.185)	4.74	4.20

59/Bd/1.3

Barley, Grain (at 85% dry matter): cwt per acre						
Green manures		N: cwt per acre (including basal)		Dung to cabbages 1952: tons per acre		
In barley for potatoes	After potatoes for barley	Ryegrass	Trefoil	0.23	0.46	None
None	Under-sown			0.23	0.46	None
(±1.06)	(±1.06)	(±1.06)	(±1.06)	(±1.06)	(±1.06)	(±0.75)
22.1	24.4	23.1	23.4	21.6	24.9	22.4
26.7	30.0	30.0	26.7	25.6	31.1	27.6
<u>Excluding plots fallow under old scheme</u>						
Green manures ploughed in						
Early						
Late						
Green manures in barley for potatoes						
None						
Undersown						
Green manures after potatoes for barley						
Trefoil						
Ryegrass						
N: cwt per acre (including basal)						
0.23						
0.46						
Mean (±0.75)						
<u>Plots fallow under old scheme</u>						
Green manures after potatoes for barley						
None	Trefoil	Ryegrass	Excluding fallow	Mean		
Fallow						
Old scheme						
20.7	26.6	25.0	24.8			
(±1.06)	(±0.75)					
N: cwt per acre (including basal)		(±2.12)		(±1.50)		
0.23	0.46	0.23	0.46	13.2	19.3	16.3
				26.2	24.0	25.1
Mean (±1.50)		19.7		21.6		
		20.7		20.7		

Mean dry-matter % as harvested: 84.4

59/Be/1.1

LEY AND ARABLE ROTATIONS

Woburn Stackyard 1959 - the 22nd year.

For history, treatments etc., see "Details of the Classical and Long Term Experiments" 1956.

Combine harvesting: In 1959 the combine harvester was introduced for barley and rye; the whole plot area was harvested for yield estimates.

Cultivations, etc.,

Treatment crops

Ley rotations

Ley 1st year. Ploughed twice: Sept 17 and Nov 21, 1958. PK fertilizers and 'Nitra-Shell' applied: Mar 31, 1959. Seed sown at 40 lb per acre: Apr 7. Sprayed with DNBP at 9 pints in 80 gallons per acre: June 4. 'Nitra-Shell' applied: 2nd dressing - Aug 31; 3rd dressing - Oct 7. Grazed 3 circuits: June 30 - Oct 30. Seeds mixture: 20 lb S24 Perennial Ryegrass, 11 lb S143 Cocksfoot, 6 lb Late Flowering Red Clover, 3 lb S100 White Clover per acre.

Ley 2nd year. Potash and nitrogen fertilizer applied: Mar 17, June 10 and Aug 31. Grazed 5 circuits: Apr 22 - Aug 29.

Ley 3rd year. Potash and nitrogen fertilizer applied: Mar 17, June 10 and Aug 31. Grazed 5 circuits: Apr 14 - Aug 21.

Lucerne 1st year. Ploughed twice: Sept 17 and Nov 21, 1958. PK fertilizer applied: Mar 31. Seed sown at 25 lb per acre: Apr 7.* Sprayed with dieldrin at 2 pints in 40 gallons per acre against birds: June 4. Cut twice: July 10, Sept 2.

Variety: Du Puits.

Lucerne 2nd year. Muriate of potash applied: Apr 13. Cut 3 times: June 12, July 10, Sept 2.

Lucerne 3rd year. Plots 3, 4, 9 and 10 were fallowed because of stem eelworm and received no potash. Ploughed: Apr 30, June 13, Sept 2, 1959.

Arable rotations

Potatoes 1st course. Ploughed twice: Sept 17 and Nov 21, 1958. Compound fertilizer applied: Mar 31, 1959. Potatoes machine planted: Apr 1. Earthed up: June 22. Sprayed with zineb at 2 lb and demeton methyl (against aphids) at 12 fluid oz (50% active ingredients) in 40 gallons per acre: Aug 15. Haulm destroyed mechanically: Sept 22. Lifted: Sept 30. Variety: Majestic.

Rye 2nd course. Ploughed: Oct 16, 1958. Seed drilled at $2\frac{1}{2}$ bushels per acre: Oct 21. 'Nitra-Shell' applied: Apr 13, 1959. Seeds hay mixture undersown on 4 plots: Apr 7. Combine harvested: Aug 18. Variety: King II.

*Sprayed with dieldrin at 2 pints in 40 gallons per acre against weevil: Apr 30.