

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/W/BG/1 Irrigation

Rothamsted Research

Rothamsted Research (1960) *59/W/BG/1 Irrigation* ; Yields Of The Field Experiments 1959, pp 61 - 66 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Bg/1.2

Cultivations, etc.:

Sugar beet. Ploughed: Oct 30 and Dec 6, 1958. Ground chalk applied: Dec 4. Salt applied: Mar 12, 1959. Basal fertilizer and 'Nitra-Shell' applied: Apr 1. Seed drilled at 10 lb per acre: Apr 3. Singled: May 19 - 22. Sprayed with dieldrin at 2 pints in 40 gallons per acre: May 26. Sprayed with demeton methyl (50% active ingredients) at 12 fluid oz in 40 gallons per acre: June 4 and June 22. Lifted: Oct 19 - 22. Variety: Klein E.

Spring wheat. Ploughed: Nov 18, 1958. Fertilizers applied: Mar 13, 1959. Seed drilled at 3 bushels per acre: Mar 17. Combine harvested: Aug 21. Variety: Peko.

Spring beans. Ploughed: Sept 15 and Dec 3, 1958. Dung applied: Dec 2. Seed combine drilled at 200 lb per acre: Feb 26, 1959. Combine harvested: Aug 7 and 22. Variety: Garton's Spring Tick.

Grass. Basal fertilizers applied: Nov 24, 1958. 'Nitra-Shell' and 0.6 cwt muriate of potash per acre applied: Mar 24, 1959. Cut six times (all plots): May 6 and 27, June 22, July 14, Aug 11, Sept 9. 'Nitra-Shell' applied after each cut, except the last. 2nd dressing of 0.6 cwt muriate of potash per acre applied after the third cut. Variety: Cocksfoot S37.

Standard errors per plot.

Sugar beet.	Total sugar, whole plot:	5.88 cwt per acre or 8.0%
		(6 d.f.)
	sub plot:	4.23 cwt per acre or 5.7%
		(8 d.f.)
Tops,	whole plot:	1.467 cwt per acre or 16.5%
		(6 d.f.)
	sub plot:	0.870 cwt per acre or 9.8%
		(8 d.f.)
Spring wheat.	Grain (at 85% D.M.),	whole plot: 2.55 cwt per acre or 11.3%
		(6 d.f.)
	sub plot:	2.74 cwt per acre or 12.1%
		(8 d.f.)
Cut grass.	Dry matter, Total of cuts 1 - 3	whole plot: 2.10 cwt per acre or 8.0%
		(6 d.f.)
	sub plot:	1.37 cwt per acre or 5.2%
		(8 d.f.)
	Total of cuts 4 - 6	whole plot: 2.23 cwt per acre or 10.1%
		(6 d.f.)
sub plot:	1.95 cwt per acre or 8.8%	
	(8 d.f.)	
Total of cuts 1 - 6	whole plot: 3.36 cwt per acre or 7.0%	
	(6 d.f.)	
sub plot:	3.18 cwt per acre or 6.6%	
	(8 d.f.)	

Summary of Results

Sugar beet

Roots washed: tons per acre

Spray	Irrigation				
	0	C			
None	14.64	21.91			
Demeton methyl	14.81	22.40			
N: cwt per acre			None	Spray Demeton methyl	Mean
0.6	14.90	21.70	18.06	18.54	18.30
1.2	14.55	22.61	18.50	18.67	18.58
Mean	14.73	22.16	18.28	18.61	18.44
Difference	-0.35	+0.91	+0.44	+0.13	+0.28

Sugar percentage

Spray	Irrigation				
	0	C			
None	20.1	19.7			
Demeton methyl	20.5	19.9			
N: cwt per acre			None	Spray Demeton methyl	Mean
0.6	20.8	20.3	20.3	20.8	20.5
1.2	19.8	19.4	19.6	19.6	19.6
Mean	20.3	19.8	19.9	20.2	20.1
Difference	-1.0	-0.9	-0.7	-1.2	-0.9

Total sugar: cwt per acre

Spray	Irrigation				
	0	C			
None	(±3.40)				
Demeton methyl	58.5	86.4			
	60.8	89.0			
N: cwt per acre			None	Spray Demeton methyl	Mean
0.6	(±2.69)*				
	61.8	87.8	72.8	76.8	74.8
1.2	57.5	87.6	72.1	73.0	72.6
Mean (±2.40)	59.7	87.7	72.5	74.9	73.7
Difference (±2.44)	-4.3	-0.2	-0.7	-3.8	-2.2 (±1.73)

* for use in horizontal and diagonal comparisons only.

59/Bg/1.4

Sugar beet
Tops: tons per acre

Spray	Irrigation		Spray		
	0	C	None	Demeton methyl	Mean
	(±0.847)				
None	7.79	10.23			
Demeton methyl	6.78	10.67			
N: cwt per acre					
	(±0.649)*		(±0.649)*		
0.6	6.65	9.05	7.91	7.79	7.85
1.2	7.92	11.84	10.11	9.66	9.88
Mean (±0.599)	7.29	10.45	9.01	8.73	8.87
Difference (±0.503)	+1.27	+2.79	+2.20	+1.87	+2.03 (±0.355)

Spring wheat

Grain (at 85% dry matter): cwt per acre

N: cwt per acre	0	Irrigation			Mean
		A	B	C	
		(±1.85)*			(±0.79)
0.4	18.2	18.9	23.9	28.7	22.4
0.8	17.0	16.4	27.8	30.0	22.8
Mean (±1.48)	17.6	17.6	25.9	29.4	22.6
Difference (±2.24)	-1.2	-2.5	+3.9	+1.3	+0.4 (±1.12)

Spring beans

Grain (at 85% dry matter): cwt per acre

Dung: tons per acre	Treatment								Mean
	0	OS	C ₁	C ₁ S	C ₂	C ₂ S	C ₃	C ₃ S	
None	10.3 ⁺	8.9 ⁺	28.5	25.6	25.3	23.6	23.0	23.1	17.2
12	11.9 ⁺	9.2 ⁺	24.6	26.3	23.5	26.4	22.4	24.8	17.6
Mean	11.1	9.1	26.6	26.0	24.4	25.0	22.7	24.0	17.4
Difference	+1.6	+0.3	-3.9	+0.7	-1.8	+2.8	-0.6	+1.7	+0.4
Mean dry matter % as harvested: 84.9									

* for use in horizontal and diagonal comparisons only.

⁺ means of 3 sub plots. All other values in body of table are based on 1 sub plot only.

Cut grass

Total of cuts 1 - 3. Dry matter: cwt per acre

K ₂ O: cwt per acre including basal	Irrigation		K ₂ O: cwt per acre including basal		Mean
	0	C			
	(±1.21)				
1.2	19.8	32.5			
1.8	16.6	35.3			
N: cwt per acre ⁺			1.2	1.8	
	(±0.94)*		← (±0.94)*		
0.3	15.4	29.3	22.6	22.1	22.4
0.6	20.9	38.5	29.6	29.8	29.8
Mean (±0.86)	18.2	33.9	26.1	26.0	26.1
Difference (±0.79)	+5.5	+9.2	+7.0	+7.7	+7.4 (±0.56)

Total of cuts 4 - 6. Dry matter: cwt per acre

K ₂ O: cwt per acre including basal	Irrigation		K ₂ O: cwt per acre including basal		Mean
	0	C			
	(±1.29)				
1.2	13.0	27.5			
2.4	13.8	34.6			
N: cwt per acre ⁺			1.2	2.4	
	(±1.07)*		← (±1.07)*		
0.3	12.4	27.8	18.6	21.6	20.1
0.6	14.5	34.3	22.0	26.8	24.4
Mean (±0.91)	13.4	31.1	20.3	24.2	22.2
Difference (±1.12)	+2.1	+6.5	+3.4	+5.2	+4.3 (±0.79)

* for use in horizontal and diagonal comparisons only.

⁺ for each cut.

Mean dry matter \bar{x} as cut:

Total of cuts 1 - 3: 23.6

Total of cuts 4 - 6: 24.7

Cut grass

Total of cuts 1 - 6. Dry matter: cwt per acre

K ₂ O: cwt per acre including basal	Irrigation		K ₂ O: cwt per acre including basal		Mean
	0	C			
	(±1.94)				
1.2	32.8	60.1			
2.4	30.4	69.9			
N: cwt per acre ⁺			1.2	2.4	
	(±1.65)*			← (±1.65)*	
0.3	27.8	57.1	41.2	43.7	42.5
0.6	35.4	72.9	51.7	56.6	54.2
Mean (±1.37)	31.6	65.0	46.4	50.2	48.3
Difference (±1.83)	+7.6	+15.8	+10.5	+12.9	+11.7 (±1.30)

* for use in horizontal and diagonal comparisons only.

⁺ for each cut.

Mean dry matter % as cut:

Total of cuts 1 - 6: 24.2

WINTER WHEAT

Seed rates, sowing dates and levels of nitrogen (after non-cereal crop) - Great Field I 1959.

Design: 3 randomized blocks of 9 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0148 acres. Area harvested: 0.0096 acres.

Treatments. All combinations of:-

Whole plots. Seed rates: 2; 3; 4 bushels per acre.
Sowing dates: Oct 16; Nov 21, 1958; Jan 8*, 1959.

Sub plots. Nitrogen (in addition to basal): 0.47; 0.93 cwt N per acre applied as 'Nitro-Chalk' in two equal parts in February and April.

*Note. Sowing on one block delayed by bad weather until Jan 24, 1959.

Basal dressing: 3 cwt compound fertilizer (10% P₂O₅, 20% K₂O) per acre broadcast in seed bed, 3 cwt compound fertilizer (5% N, 12½% P₂O₅, 12½% K₂O) per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Sept 12, 1958. Compound fertilizer applied: First sowing - Oct 16; second sowing - Nov 21; third sowing - Jan 8, 1959 (plots 2, 4 and 5 - Jan 26). First dressing of N applied: Feb 16. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: Apr 21. Second dressing of N applied: Apr 22. Combine harvested: Aug 17. Variety: Cappelle. Previous crop: Potatoes.

Note. Counts of plant shoot and ear number, and estimates of plant height and % area lodged were made. Severe lodging occurred in early July and the mean % areas lodged at harvest were:

Sowing date	%	Seed rate		N	
		bu. p.a.	%	c.p.a.	%
Oct 16	94	2	46	0.6	42
Nov 21	66	3	57	1.1	73
Jan 8	14	4	70		

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 2.48 cwt per acre or 4.9% (16 d.f.)
Sub plot: 3.38 cwt per acre or 6.7% (18 d.f.)

Errata to 'Results of the Field Experiments' 1958 page 58/Ca/1.2

Rates of N cwt per acre should read '0.6' and '1.2' not '0.4' and '0.8'.
S.E. of means of seed rates and sowing dates should read '0.79' not '0.56'.