

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/R/CA/3 Spring Wheat - Row Spacing, Seed Rates and N

Rothamsted Research

Rothamsted Research (1960) *59/R/CA/3 Spring Wheat - Row Spacing, Seed Rates and N* ; Yields Of The Field Experiments 1959, pp 71 - 72 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Ca/3.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Row spacing: inches	7	7	14	14	7B	
Seed rate: bushels per acre	2	4	1	2	3	Mean
Mean (± 0.68)	33.0	33.6	29.9	30.0	31.8	31.7
<u>Drill</u>						
Standard (± 0.95)	33.1	32.7	29.6	27.7	33.1	31.2
Precision	33.0	34.5	30.3	32.4	30.5	32.1
Difference (± 1.35)	-0.1	+1.8	+0.7	+4.7	-2.6	± 0.9 (± 0.60)
<u>N cwt per acre</u>						
0.6 (± 1.16)*	31.2	31.7	29.0	29.6	31.9	30.7
1.2	34.9	35.5	30.8	30.5	31.7	32.7
Difference (± 1.90)	+3.7	+3.8	+1.8	+0.9	-0.2	+2.0 (± 0.85)

	Standard	Drill Precision	Difference
		(± 0.74)**	(± 1.04)
<u>N cwt per acre</u>			
0.6	30.6	30.7	+0.1
1.2	31.9	33.5	+1.6
Difference (± 1.20)	+1.3	+2.8	+1.5 (± 1.69)

* For use only in horizontal and diagonal comparisons.

** For use only in diagonal comparisons.

B = every 4th row blank.

SPRING WHEAT

Combine drilling of nitrogen - Rothamsted (R) Deacons Field and Woburn (W) Lansome Field 1959.

Design (each field): 4 randomized blocks of 7 plots each.

Area of each plot:

Deacons Field (R): 0.0212 acres. Area harvested: 0.0141 acres.
Lansome Field (W): 0.0186 acres. Area harvested: 0.0124 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.2; 0.5; 0.8 cwt N per acre.

Method of application: Broadcast as sulphate of ammonia; combine drilled as compound fertilizer:

N₁: 6% N, 15% P₂O₅, 15% K₂O.

N₂: 8% N, 8% P₂O₅, 8% K₂O.

N₃: 12% N, 9% P₂O₅, 9% K₂O.

Basal dressing per acre (each field): 0.54 cwt P₂O₅ and 0.54 cwt K₂O combine drilled

(a) as compound 16% P₂O₅, 16% K₂O on the no nitrogen and broadcast nitrogen plots;

(b) as compounds N₁, N₂, N₃ on the plots receiving drilled nitrogen.

Cultivations, etc.:

Deacons Field (R): Ploughed: Nov 17, 1958. Seed combine drilled at 3 bushels per acre; balance of compound fertilizer (16% P₂O₅, 16% K₂O) and sulphate of ammonia applied: Mar 19, 1959. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 6. Combine harvested: Aug 21. Variety: Koga II. Previous crop: Winter wheat.

Lansome Field (W): Ploughed: Nov 5, 1958. Seed combine drilled at 3 $\frac{1}{3}$ bushels per acre: Mar 21, 1959. Sulphate of ammonia applied: Mar 23. Combine harvested: Aug 21. Variety: Peko. Previous crop: Spring Wheat.

Standard errors per plot, Grain (at 85% dry matter):

Deacons Field (R): 0.90 cwt per acre or 3.8% (18 d.f.)

Lansome Field (W): 1.34 cwt per acre or 5.9% (18 d.f.)

Note: Plant counts at germination were made.