

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1959

[Full Table of Content](#)



59/W/CA/5 Wheat - Time of Sowing, Forms of N

Rothamsted Research

Rothamsted Research (1960) *59/W/CA/5 Wheat - Time of Sowing, Forms of N* ; Yields Of The Field Experiments 1959, pp 75 - 76 - DOI: <https://doi.org/10.23637/ERADOC-1-179>

59/Ca/5.2

Summary of Results

Time of sowing	N: fertilizer applied					Mean
	None	Autumn		Spring		
		C	S	C	S	
<u>Grain (at 85% dry matter): cwt per acre</u>						
			(± 2.04)*			
Winter	11.5	16.7	20.4	19.4	24.8	18.6
Spring	13.7	10.8	13.2	15.4	11.5	12.9
Mean (± 1.44)	12.6	13.8	16.8	17.4	18.1	15.7
Diff. (± 2.88)**	+2.2	-5.9	-7.2	-4.0	-13.3	-5.7

* For use in horizontal comparisons only.

** For use only in testing the difference of two differences.

Mean dry matter % as harvested: 86.2

Straw (at 85% dry matter): cwt per acre

Winter	10.3	12.3	18.7	23.4	24.1	17.7
Spring	13.3	12.3	16.0	26.9	22.1	18.1
Mean	11.8	12.3	17.3	25.1	23.1	17.9
Diff.	+3.0	0.0	-2.7	+3.5	-2.0	+0.4

Mean dry matter % as harvested: 76.3

C = 100 lb N per acre as calcium nitrate

S = 100 lb N per acre as sulphate of ammonia.

59/Cb/t.1

BARLEY

Combine drilling of nitrogen - Rothamsted (R) Deacons Field and Woburn (W) Lansome Field 1959.

Design: 4 randomized blocks of 7 plots each.

Area of each plot:

Deacons Field (R): 0.0212 acres. Area harvested: 0.0141 acres.
Lansome Field (W): 0.0186 acres. Area harvested: 0.0124 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.2; 0.5; 0.8 (0.7 - Lansome Field) cwt N per acre.

Methods of application: Broadcast as sulphate of ammonia; combine drilled as compound fertilizer:

N_1 : 6% N; 15% P_2O_5 ; 15% K_2O .
 N_2 : 8% N; 8% P_2O_5 ; 8% K_2O .
 N_3 : 12% N; 9% P_2O_5 ; 9% K_2O .

Basal dressing per acre:-

Deacons Field (R): 0.57 cwt P_2O_5 and 0.57 cwt K_2O
Lansome Field (W): 0.54 cwt P_2O_5 and 0.54 cwt K_2O .

combine drilled

(a) as compound (16% P_2O_5 , 16% K_2O) on the no nitrogen and broadcast nitrogen plots,

(b) as compounds N_1 , N_2 , N_3 on the plots receiving drilled nitrogen.

Cultivations, etc.:

Deacons Field (R): Ploughed: Nov 17-24, 1958. Seed combine drilled at 2 bushels per acre: Mar 19, 1959. Sulphate of ammonia and balance of compound fertilizer (16% P_2O_5 , 16% K_2O) broadcast: Mar 20. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 6. Combine harvested: Aug 6. Variety: Proctor. Previous crop: Winter wheat.

Lansome Field (W): Ploughed: Nov 5, 1958. Seed combine drilled at 3 bushels per acre: Mar 20, 1959. Sulphate of ammonia broadcast: Mar 23. Combine harvested: Aug 17. Variety: Herta. Previous crop: Barley.

Standard errors per plot, Grain (at 85% dry matter):

Deacons Field (R): 1.69 cwt per acre or 6.0% (18 d.f.)
Lansome Field (W): 1.70 cwt per acre or 6.4% (18 d.f.)

Note: Plant counts at germination were made.