

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/C/8 Intensive Barley Growing - Cereals and Beans

Rothamsted Research

Rothamsted Research (1962) *61/R/C/8 Intensive Barley Growing - Cereals and Beans* ; Yields Of The Field Experiments 1961, pp 105 - 106 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/C/8.1

INTENSIVE BARLEY GROWING EXPERIMENT

Little Knott I - 1961

Design: 2 replicates of 40 treatments in 4 blocks of 20 plots each.

Area of each plot (acres): 0.0212. Area harvested: 0.0138

Treatments. All combinations of:-

Crop sequences:

	1961	1962	1963	1964	1965	1966	1967	1968
1	O	Be	B	B	B	B	B	B
2	SW	O	Be	B	B	B	B	B
3	O	SW	O	Be	B	B	B	B
4	Be	O	SW	O	Be	B	B	B
5	SW	Be	O	SW	O	Be	B	B
6	SW	SW	Be	O	SW	O	Be	B
7	B	B	B	B	B	B	B	B
8	SW	SW	SW	SW	SW	SW	SW	SW
9	W*	W	W	W	W	W	W	W
10	Be	W	P	B	Be	W	P	B

O = Oats, Be = Spring beans, SW = Spring wheat, W = Winter wheat,
B = Barley, P = Potatoes.

Nitrogen: Applied to continuous crops and to the winter wheat and
barley in treatment 10 - none; 0.3; 0.6; 0.9 cwt N per acre
as 'Nitro-Chalk'.

*In this case, because of bad weather, spring instead of winter wheat
was sown.

Basal dressings (per acre): 240 lb compound fertiliser, 14% P₂O₅,
28% K₂O, to all crops except potatoes, which receive 10 cwt
compound fertiliser, 10% N, 10% P₂O₅, 18% K₂O. The non-continuous
crops oats and spring wheat also receive 0.45 cwt N as 'Nitro-
Chalk'.

Cultivations, etc.: Ground chalk applied at 24 cwt per acre: Sept 23,
1960. Ploughed: Oct 4.

Spring beans: Seed placement drilled at 200 lb per acre: Mar 10, 1961.
Sprayed with demeton-methyl at 12 fluid oz in 60 gallons per acre:
June 12. Combine harvested: Aug 26. Variety: Tick.

Oats: Seed combine drilled at 4 bushels per acre, 'Nitro-Chalk'
applied: Mar 10, 1961. Sprayed with MCPA/TBA at 4 pints in
40 gallons per acre: May 12. Combine harvested: Aug 22.
Variety: Condor.

Spring wheat: Seed combine drilled at 3 bushels per acre, 'Nitro-
Chalk' applied: Mar 10, 1961. Sprayed with MCPA/TBA at
4 pints in 40 gallons per acre: May 12. Combine harvested:
Aug 30. Variety: Jufy I.

Barley: Seed combine drilled at 2½ bushels per acre: Mar 9, 1961.
'Nitro-Chalk' applied: Mar 10. Sprayed with MCPA/TBA at 4
pints in 40 gallons per acre: May 12. Combine harvested:
Aug 30. Variety: Proctor.

Previous crop: Spring wheat.

61/C/8.2

Note. Yields were only taken for sequences 2, 7, 8 and 9.

Standard error per plot. Grain (at 85% dry matter):
 Spring wheat (8 & 9): 2.27 cwt per acre or 8.2% (11 d.f.)
 Spring wheat (2): 2.70 cwt per acre or 9.2% (6 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Spring wheat (8 and 9)

N: cwt per acre

None	0.3	0.6	0.9	Mean
16.9	25.3	32.0	36.0	27.5
(±1.13)				
Mean dry matter % as harvested: 85.5				

Spring wheat (2)

N: cwt per acre

0.45

29.3

(±0.96)

Mean dry matter % as harvested: 85.8

Barley (7)

N: cwt per acre

None	0.3	0.6	0.9	Mean
22.7	34.5	38.8	38.8	33.7

Mean dry matter % as harvested: 83.9