


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/DA/4 Wheat - Sowing Dates, Seed Rates and N (After Non-cereal Crop)

Rothamsted Research

Rothamsted Research (1962) *61/R/DA/4 Wheat - Sowing Dates, Seed Rates and N (After Non-cereal Crop)* ; Yields Of The Field Experiments 1961, pp 111 - 112 - DOI:

<https://doi.org/10.23637/ERADOC-1-182>

61/Da/4.1

WINTER AND SPRING WHEAT

Sowing dates, seed rates and levels of nitrogen (after non-cereal crop) - Great Knott III 1961.

Design: 3 randomised blocks of 12 plots each, plots split into 2 for the application of nitrogen.

Area of each sub plot: 0.0148 acres. Area harvested: 0.0097 acres.

Treatments. All combinations of:-

Whole plots. Sowing dates: Sept 29, 1960; Jan 19, 1961;
Feb 15 (winter wheat); Feb 15* (spring wheat).
Seed rates: 2; 3; 4 bushels per acre.

Sub plots. Nitrogen (in addition to basal): 0.46; 0.92 cwt
N per acre applied as 'Nitro-Chalk' in two equal parts on
February 16 and April 19.

* Weather conditions delayed sowing and spring wheat was drilled on Feb 15 instead of a still later sowing of winter wheat.

Basal dressing: 2 cwt compound fertiliser (14% P_2O_5 , 28% K_2O) per acre broadcast in seedbed, 3 cwt compound fertiliser (5% N, 12½% P_2O_5 , 12½% K_2O) per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Sept 19, 1960. Compound fertiliser applied: First sowing - Sept 29; second sowing - Nov 16; third sowing - Feb 15, 1961. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 5. Combine harvested: Aug 30. Varieties: Cappelle and Jufy I. Previous crops: 1958 - Spring wheat; 1959 - Winter and spring beans; 1960 - Early potatoes.

Note. Counts of plant shoot and ear number, and estimates of plant height and % area lodged were made.

Standard error per plot, Grain (at 85% dry matter):

Whole plot: 3.30 cwt per acre or 7.4% (22 d.f.)

Sub plot: 3.30 cwt per acre or 7.4% (24 d.f.)

61/Da/4.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Seed rate: bushels per acre	Variety and date of sowing				N: cwt per acre (including basal)		Diff.	Mean	
	Cappelle (W.W.) Sept 29th	Jan 19th	Feb 15th	July I (S.W.) Feb 15th	0.6	1.1			
	(±1.91)				(±1.17)*		(±1.35)	(±0.95)	
2	44.0	49.0	43.7	41.3	43.3	45.6	+2.3	44.5	
3	39.9	47.7	45.7	43.7	43.5	45.0	+1.5	44.2	
4	39.7	49.0	47.9	44.9	45.3	45.4	+0.1	45.4	
	Variety and date of sowing				(±1.35)*		(±1.56)	(±1.10)	
	Cappelle (W.W.)				Sept 29th	42.9	39.5	-3.4	41.2
	Cappelle (W.W.)				Jan 19th	46.3	50.7	+4.4	48.5
	Cappelle (W.W.)				Feb 15th	45.7	45.8	+0.1	45.7
	July I (S.W.)				Feb 15th	41.3	45.3	+4.0	43.3
	Mean					44.0	45.4	+1.4	44.7
								(±0.78)	

*For use in vertical and diagonal comparisons only.

Mean dry matter % as harvested: 86.6