

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1961

[Full Table of Content](#)



---

## 61/R/DB/1 Barley - Forms and Application of N

### Rothamsted Research

Rothamsted Research (1962) *61/R/DB/1 Barley - Forms and Application of N* ; Yields Of The Field Experiments 1961, pp 115 - 116 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/Db/1.1

BARLEY

Forms and methods of application of nitrogen - Great Knott I 1961.

Design: 3 randomised blocks of 20 plots each.

Area of each plot: 0.0194 acres.

Treatments: No nitrogen (2 plots per block) and all combinations of:-

Forms of N: Ammonium sulphate 21% N (S);  
Calcium nitrate 15.5% N (C);  
Urea 45.6% N (U)

Levels of N: 0.35; 0.70 cwt N per acre.

Methods of application: Broadcast (B); combine drilled (D);  
side band placed (P).

Basal dressing: 2 cwt granular compound fertiliser (14%  $P_2O_5$ , 28%  $K_2O$ )  
per acre.

Cultivations, etc.: Ground chalk applied at 45 cwt per acre:

Oct 17, 1960. Ploughed: Nov 29. Seed drilled at  $2\frac{1}{2}$  bushels  
per acre and seedbed fertilisers applied: Mar 14, 1961. Sprayed  
with CMFP at 6 pints in 40 gallons per acre: May 11. Combine  
harvested: Aug 16. Variety: Proctor. Previous crop: Winter wheat.

Standard error per plot.

Grain (at 85% dry matter): 1.61 cwt per acre or 4.1% (37 d.f.)

61/Db/1.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Form of N			Mean
	S	C	U	
Mean ( $\pm 0.38$ )	39.2	39.6	38.1	39.0 ( $\pm 0.22$ )
N: cwt per acre				
0.35 ( $\pm 0.54$ )	36.4	36.3	36.0	36.2
0.75	42.1	42.8	40.2	41.7
Diff. ( $\pm 0.76$ )	+5.7	+6.5	+4.2	+5.5 ( $\pm 0.44$ )
Method of application				
B	38.0	41.1	38.0	39.0
D ( $\pm 0.66$ )	40.3	38.3	37.1	38.5 ( $\pm 0.38$ )
P	39.5	39.4	39.2	39.3
	Method of application			
	B	D	P	
N: cwt per acre				
0.35 ( $\pm 0.54$ )	35.8	36.5	36.3	
0.70	42.2	40.6	42.3	
Diff. ( $\pm 0.76$ )	+6.4	+4.1	+6.0	

No N: 25.0 ( $\pm 0.66$ )

General mean: 37.6

Mean dry matter % as harvested (all plots): 82.5

Form of N

S = Ammonium sulphate 21% N

C = Calcium nitrate 15.5% N

U = Urea 45.6% N

Method of application

B = Broadcast

D = Combine drilled

P = Side band placed