


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1961

[Full Table of Content](#)



---

## 61/R/DG/1 Grass - Levels of K and N 4th Year

### Rothamsted Research

Rothamsted Research (1962) *61/R/DG/1 Grass - Levels of K and N 4th Year* ; Yields Of The Field Experiments 1961, pp 135 - 136 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/Dg/1.1

## GRASS

Levels of N and K - Harwoods Piece 1961 - the 4th year.

Design: 4 randomised blocks of 12 plots each.

Area of each plot: 0.0087 acres. Area harvested: 0.0057 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre as 'Nitro-Chalk'.

Potash: None; 0.3; 0.6 cwt  $K_2O$  per acre as muriate of potash.

All the above in the presence of 0.6 cwt  $P_2O_5$  per acre as superphosphate.

In addition 2 plots per block, receiving 0.9 cwt N and 0.6 cwt  $K_2O$  per acre, also received phosphate at either None or 1.2 cwt  $P_2O_5$  per acre as superphosphate.

Note: (1) N and K dressings are applied for each cut. All P dressings are applied once annually.

(2) All treatments were applied to the same plots as in the previous seasons.

Basal dressing: None.

Cultivations, etc.: Sprayed with dalapon at 8 lb in 40 gallons per acre: Sept 7, 1960. 1st dressing of fertilisers applied: Mar 3, 1961. Cut twice: May 12 and July 4.

Standard errors per plot. Dry matter:

1st cut: 3.04 cwt per acre or 6.4% (33 d.f.)

2nd cut: 1.16 cwt per acre or 8.8% (33 d.f.)

Total of 2 cuts: 3.28 cwt per acre or 5.4% (33 d.f.)

Note: (3) For details of the previous years results see 'Results of the Field Experiments' 58/Cg/2, 59/Cg/2 and 60/Ci/1.

61/Dg/1.2

Summary of Results

Dry matter: cwt per acre

cwt per acre														
N*	0.0	0.3	0.3	0.3	0.6	0.6	0.6	0.9	0.9	0.9	0.9	0.9	0.9	
P <sub>20</sub> <sup>5</sup>	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.0	1.2	
K <sub>20</sub> <sup>5</sup> *	0.0	0.0	0.3	0.6	0.0	0.3	0.6	0.0	0.3	0.6	0.6	0.6	0.6	Mean
1st cut (±1.51)	12.0	43.5	39.7	39.7	53.0	54.8	55.1	52.9	52.6	53.6	54.3	54.3	47.1	
2nd cut (±0.58)	5.1	10.5	10.9	10.9	13.5	11.7	12.5	15.0	16.4	17.7	17.3	17.8	13.3	
Total of 2 cuts (±1.64)	17.2	54.0	50.6	50.6	66.5	66.5	67.6	67.9	69.0	71.3	71.6	72.1	60.4	

\*For each cut.

Mean dry matter % as cut:

1st cut: 22.1  
 2nd cut: 32.4  
 Total of 2 cuts: 27.2