


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/B/7 Concentrated Fertiliser Rotation

Rothamsted Research

Rothamsted Research (1962) *61/R/B/7 Concentrated Fertiliser Rotation* ; Yields Of The Field Experiments 1961, pp 74 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/B/7.1

CONCENTRATED FERTILISER ROTATION

Concentrated compound fertiliser and forms of N - West Barnfield I
1961, the second year.

Rotation: Kale, ryegrass, barley.

Design (each crop): 2 randomised blocks of 14 plots each.

Area of each plot (acres): 0.0174. Area harvested: Kale -
0.0084, ryegrass - 0.0061, barley - 0.0116.

Treatments (per acre): No fertiliser (O)
 P_2O_5 and K_2O each at 0.3 cwt to barley and each at 1.0 cwt to
kale and ryegrass, as triple superphosphate and potassium
bicarbonate. (B)
Compound fertiliser, 20% N, 10% P_2O_5 , 10% K_2O at 0.3(1),
0.6(2) cwt N to barley and 1.0(1), 2.0(2) to kale and
ryegrass. (F)
Sulphate of ammonia, granular superphosphate and muriate of
potash at rates equivalent to treatments F(1) and (2). (P)
PK as treatment B plus
Sulphate of ammonia (S)
Calcium nitrate (C)
Urea (U)
Ammonium nitrate (A)
each at rates 1 and 2 of N.

Basal dressing: None.

Cultivations, etc.: Ploughed: Jan 11, 1961. Fertilisers broadcast for
barley, barley drilled at $2\frac{1}{2}$ bushels per acre: Mar 13. Fertilisers
broadcast for ryegrass, ryegrass sown at 30 lb per acre: Mar 17.
Fertilisers applied for kale: Apr 11. Kale drilled at 3 lb per
acre: Apr 24. Barley and ryegrass sprayed with CMPP at 6 pints
in 40 gallons per acre: May 13. Grass cut: July 20. Barley
combine harvested: Aug 16. Grass cut second time: Sept 26.
Kale harvested: Nov 6 and 16. Varieties: Kale - Thousand head
(Canson); ryegrass - S22; barley - Proctor.

Erratum to "Results of the Field Experiments" 1960 page 60/B/8.1(q.v.):
Treatment B should read " P_2O_5 and K_2O each at 0.3 cwt to barley and each
at 1.0 cwt to kale and ryegrass" and not as shown.

Standard errors per plot.

Kale, fresh weight: 1.343 tons per acre or 6.6% (13 d.f.)

Ryegrass dry matter:

1st cut: 3.10 cwt per acre or 4.5% (13 d.f.)

2nd cut: 1.21 cwt per acre or 13.2% (13 d.f.)

Total of 2 cuts: 3.43 cwt per acre or 11.2% (13 d.f.)

Barley, grain (at 85% dry matter): 1.66 cwt per acre or 5.9% (13 d.f.)

61/B/7.2

Summary of Results

Fertiliser	Kale fresh weight: tons per acre	Ryegrass dry matter: cwt per acre			Barley (at 85% dry matter): cwt per acre	
		1st cut	2nd cut	Total of 2 cuts	Grain	Straw
	(±0.949)	(±2.19)	(±0.85)	(±2.43)	(±1.18)	
O	10.91	8.2	5.5	13.7	9.3	4.5
B	13.05	7.4	5.0	12.3	12.6	5.2
F ₁	19.48	21.5	9.5	31.0	27.1	16.0
F ₂	23.72	26.4	11.4	37.7	37.0	27.9
P ₁	20.14	19.9	8.8	28.6	25.5	15.1
P ₂	23.91	24.2	10.4	34.6	37.3	28.4
S ₁	20.96	20.6	7.3	27.9	24.1	14.0
S ₂	21.97	26.4	13.2	39.5	34.3	24.7
C ₁	21.39	24.3	8.6	32.9	27.0	16.3
C ₂	23.64	26.2	10.4	36.5	36.4	22.8
U ₁	19.46	20.2	7.9	28.1	26.4	16.3
U ₂	22.60	23.5	10.8	34.3	35.0	26.4
A ₁	19.42	25.2	9.5	34.6	27.1	18.4
A ₂	22.84	24.8	10.5	35.3	34.2	21.8
Mean	20.25	21.3	9.2	30.5	28.1	18.4
Mean dry matter % as harvested:		25.0	22.2	23.6	80.8	78.1

Treatments

- O = No fertiliser
- B = P₂O₅ and K₂O each at 0.3 cwt to barley and each at 1.0 to kale and ryegrass, as triple superphosphate and potassium bicarbonate.
- F = Compound fertiliser, 20% N, 10% P₂O₅, 10% K₂O at 0.3(1), 0.6(2) cwt N to barley and 1.0(1), 2.0(2) to kale and ryegrass.
- P = Sulphate of ammonia, granular superphosphate and muriate of potash at rates equivalent to treatments F (1) and (2).
- S = Sulphate of ammonia Plus PK as treatment B
- C = Calcium nitrate " " " " "
- U = Urea " " " " "
- A = Ammonium nitrate " " " " "