


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/B/9 N Levels and Residues Rotation

Rothamsted Research

Rothamsted Research (1962) *61/R/B/9 N Levels and Residues Rotation* ; Yields Of The Field Experiments 1961, pp 79 - 80 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/B/9.1

N LEVELS AND RESIDUES ROTATION

Direct and residual effects of sulphate of ammonia - Long Hoos III 1961, the 2nd year.

Rotation: Wheat, potatoes.

Design (each crop): 3 randomised blocks of 9 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

Nitrogen (applied as sulphate of ammonia) at 3 levels in 1960 and at 3 levels in 1961.

To Wheat: None; 0.5; 1.0 cwt N per acre

To Potatoes: None; 0.75; 1.50 cwt N per acre.

Basal dressing (per acre):

To wheat: $2\frac{1}{4}$ cwt compound fertiliser, 14% P_2O_5 , 28% K_2O combine drilled.

To potatoes: 5 cwt compound fertiliser, 14% P_2O_5 , 28% K_2O broadcast on the flat.

Cultivations, etc.:

Wheat: Ground chalk applied at 23 cwt per acre, ploughed:

Mar 7, 1961. Rotary cultivated: Mar 20. Sulphate of ammonia applied, seed combine drilled at 3 bushels per acre: Mar 21.

Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 19.

Combine harvested: Aug 30. Variety: Jufy I.

Potatoes: Ground chalk applied at 23 cwt per acre: Dec 16, 1960.

Ploughed: Dec 29. Basal fertiliser broadcast on flat:

Apr 17, 1961. Rotary cultivated, sulphate of ammonia applied, ridged, potatoes planted: Apr 24. Earthed up: July 6.

Sprayed with zineb at 2 lb in 40 gallons per acre: Aug 3.

Sprayed with undiluted BOV at 15 gallons per acre: Sept 21.

Lifted: Oct 5. Variety: Ulster Supreme.

Standard errors per plot.

Wheat. Grain (at 85% dry matter): 2.18 cwt per acre or 6.5%
(16 d.f.)

Potatoes. Total Tubers: 0.775 tons per acre or 8.4% (16 d.f.)

61/B/9.2

Summary of Results

Wheat, Grain (at 85% dry matter): cwt per acre

	N: cwt per acre in 1961			Mean
	None	0.5	1.0	
N: cwt per acre to potatoes in 1960		(±1.25)		(±0.73)
None	23.8	34.9	37.9	32.2
0.75	26.1	34.2	39.0	33.1
1.50	31.0	34.5	39.2	34.9
Mean (±0.73)	27.0	34.5	38.7	33.3

Mean dry matter % as harvested: 84.9

Potatoes

	N: cwt per acre in 1961			Mean
	None	0.75	1.50	
N: cwt per acre to wheat in 1960	<u>Total tubers: tons per acre</u>			
		(±0.448)		(±0.258)
None	5.83	9.26	11.70	8.93
0.5	5.73	10.38	11.29	9.13
1.0	6.74	10.00	12.01	9.58
Mean (±0.258)	6.10	9.88	11.67	9.21

N: cwt per acre to wheat in 1960 Percentage ware (1½" riddle)

None	93.9	95.4	97.2	95.5
0.5	94.2	96.3	96.9	95.8
1.0	93.2	96.0	97.6	95.6
Mean	93.8	95.9	97.2	95.6