


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/W/B/11 Weedkiller-cultivation Rotation

Rothamsted Research

Rothamsted Research (1962) *61/W/B/11 Weedkiller-cultivation Rotation* ; Yields Of The Field Experiments 1961, pp 85 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/B/11.1

WEEDKILLER CULTIVATION ROTATION

A comparison of weed control by various cultivation methods and by a pre-emergence weedkiller - Woburn Great Hill I and II 1961.

For previous history see "Results of the Field Experiments" 60/B/11.

Rotation (commencing 1960): Potatoes, barley.

Design: 2 randomised blocks of 9 plots each per crop.

Area of each plot (acres): 0.0482. Area harvested: Potatoes - 0.0107, barley - 0.0115.

Treatments:

Potatoes: All combinations of:-

Cultivations before planting: Ploughed and spring-tine cultivated (P) - duplicate plots. Ploughed and rotary cultivated (R).

Treatments after planting: Normal cultivations (N); simazine* applied after planting (Sx); simazine* applied after early cultivations (Sy).

Barley: All combinations of:-

Cultivations: Ploughed (P); rotary cultivated (R); rigid-tine cultivated (T).

Simazine to potatoes 1960: None; 1; 2 lb per acre.

*Simazine at 1 lb active material in 40 gallons per acre.

Basal dressings per acre:

Potatoes: 10 cwt compound fertiliser, 17% N, 11% P₂O₅, 22% K₂O.

Barley: 4 cwt compound fertiliser, 16% N, 9% P₂O₅, 9% K₂O, combine drilled.

Cultivations, etc.

Potatoes: All plots ploughed: Dec 12, 1960. P plots spring-tine cultivated twice: Mar 16, 1961. P plots rolled and then spring-tine cultivated twice: Mar 23. R plots rotary cultivated: Mar 23.

Basal dressing applied, potatoes machine planted: Mar 24.

Simazine applied on rolled bouts to Sx plots: Mar 31. N and Sy plots earthed up: May 2. Sy plots earthed up and sprayed with simazine: May 19. Sx plots grubbed twice: May 30. N and Sx⁺ plots earthed up: June 14. Sprayed with undiluted BOV at 12 gallons per acre: Sept 18. Lifted: Sept 21.

Variety: Majestic.

⁺Because of failure of Sx treatment.

61/B/11.2

Barley: P plots ploughed, T plots rigid-tine cultivated twice: Feb 24, 1961. P and T plots disced: Mar 17. P and T plots rolled and spring-tine harrowed: Mar 23. R plots rotary cultivated: Mar 23. Seed combine drilled at $2\frac{1}{4}$ bushels per acre: Mar 24. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 10. Combine harvested: Aug 17. Variety: Proctor.

Standard errors per plot.

Potatoes, total tubers: 0.977 tons per acre or 16.5% (11 d.f.)
 Barley, grain (at 85% dry matter): 1.76 cwt per acre or 8.2% (8 d.f.)

Summary of Results

Potatoes

Cultivation before planting

Treatment after planting	P	R	Mean
<u>Total tubers: tons per acre</u>			
	(±0.488)	(±0.691)	(±0.399)
N	7.16	7.54	7.28
Sx	1.82	2.61	2.08
Sy	8.61	8.14	8.45
Mean	5.86 (±0.282)	6.09 (±0.399)	5.93
<u>Percentage ware ($1\frac{1}{2}$" riddle)</u>			
N	90.7	89.1	90.1
Sx	77.8	81.1	78.9
Sy	91.8	88.4	90.6
Mean	86.8	86.2	86.5

61/B/11.3

Barley

Grain (at 85% dry matter): cwt per acre

Simazine to potatoes 1960 lb per acre	Cultivation			Mean
	P	R	T	
		(±1.25)		(±0.72)
None	21.0	22.8	20.9	21.5
1	23.4	22.3	19.9	21.8
2	21.1	22.2	21.0	21.4
Mean (±0.72)	21.8	22.4	20.6	21.6