

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1961

[Full Table of Content](#)



61/R/C/4 Straw and Ryegrass Green Manure - Barley (Test Crop)

Rothamsted Research

Rothamsted Research (1962) *61/R/C/4 Straw and Ryegrass Green Manure - Barley (Test Crop)* ;
Yields Of The Field Experiments 1961, pp 97 - 97 - DOI: <https://doi.org/10.23637/ERADOC-1-182>

61/C/4

BARLEY

Effects of green manures, N and straw - Stackyard (the second year) 1961.

Design: 6 randomised blocks of 9 plots each.

Area of each plot: 0.0114 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

Nitrogen: None; 0.3; 0.6 cwt N per acre as 'Nitro-Chalk'.

Green manures and straw: None; ryegrass undersown; ryegrass undersown plus straw left on the plot after harvest. (0;R;RS)

Basal dressing: 2½ cwt per acre compound fertiliser (20% P₂O₅, 20% K₂O) combine drilled.

Cultivations, etc.: Straw spread: Sept 8, 1960. Ploughed: Dec 17.
Seed combine drilled at 2½ bushels per acre, N applied:
Mar 11, 1961. Ryegrass drilled at 40 lb per acre: Apr 18.
Sprayed with 2,4-D at ¾ pint in 40 gallons per acre: May 13.
Combine harvested: Aug 17. Variety: Proctor; Ryegrass - S22 Italian.

Standard error per plot.

Grain (at 85% dry matter): 2.31 cwt per acre or 7.7% (40 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Green manure and straw	N: cwt per acre			Mean
	None	0.3	0.6	
		(±0.94)		(±0.54)
O	21.5	33.5	40.8	31.9
R	17.6	29.4	36.8	27.9
RS	19.6	30.8	39.2	29.9
Mean (±0.54)	19.6	31.2	38.9	29.9

Mean dry matter % as harvested: 84.6

Note: For details of the previous year's results see "Results of the Field Experiments" 60/Cb/2.