


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1962

[Full Table of Content](#)



---

### 62/R/C/4 Straw and Ryegrass Green Manure - Barley (Test Crop)

#### Rothamsted Research

Rothamsted Research (1963) *62/R/C/4 Straw and Ryegrass Green Manure - Barley (Test Crop)* ;  
Yields Of The Field Experiments 1962, pp 107 - 108 - DOI:

<https://doi.org/10.23637/ERADOC-1-164>

62/C/4.1

## BARLEY

Effects of green manures, N and straw - Stackyard 1962, the 3rd year.

Design: 2 replicates of a  $3 \times 3 \times 3$  in 6 randomised blocks of 9 plots each, certain high order interactions being confounded with block differences.

Area of each plot: 0.0212 acres. Area harvested: 0.0140 acres.

Treatments. All combinations of:-

Nitrogen to barley 1962: None; 0.3; 0.6 cwt N per acre applied as 'Nitro-Chalk'.

Nitrogen to barley 1960 and 1961: None; 0.3; 0.6 cwt N per acre applied as 'Nitro-Chalk'.

Green manures and straw: None; ryegrass undersown 1960 and 1961; ryegrass undersown 1960 and 1961 plus straw left on the plot after harvest 1960 and 1961. (0; R; RS)

Basal dressing: 2 cwt compound fertiliser (20%  $P_2O_5$ , 20%  $K_2O$ ) per acre combine drilled.

Cultivations, etc.: Straw spread on "S" plots: 21 Aug, 1961. All plots ploughed: 14 Dec. 'Nitro-Chalk' applied: 21 Feb, 1962. Seed combine drilled at  $2\frac{1}{2}$  bushels per acre with basal fertiliser: 22 Feb. Sprayed with 2,4-DP/MCPA at 8 pints in 40 gallons per acre: 24 May. Combine harvested: 27 Aug. Variety: Proctor.

Note: (1) For the previous years' results see "Results of the Field Experiments" 60/Cb/2 and 61/C/4.

(2) Samples were taken for the estimation of the incidence of Take-all (Ophiobolus graminus)

Standard error per plot.

Grain (at 85% dry matter): 1.40 cwt per acre or 4.5% (22 d.f.)

Erratum to "Numerical Results of Field Experiments" 1961 page 61/C/4.1.

The area of each plot should be 0.0212 acres and not 0.0114 as given.

62/C/4.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Green manure and straw 1960 & 1961	N: cwt per acre 1962			Green manure and straw 1960 & 1961			Mean
	None	0.3	0.6	0	R	RS	
	(±0.57)						
0	21.9	32.4	38.0				
R	22.5	32.7	37.9				
RS	22.3	31.9	38.4				
N: cwt per acre 1960 & 1961	(±0.57)			(±0.57)			(±0.33)
None	21.8	31.7	37.8	30.0	31.5	29.7	30.4
0.3	21.8	32.3	37.8	30.4	30.5	31.0	30.6
0.6	23.2	33.0	38.7	31.9	31.1	31.8	31.6
Mean (±0.33)	22.2	32.3	38.1	30.8	31.0	30.8	30.9

Mean dry matter % as harvested: 81.2

Green manure and straw

- 0 = None
- R = Ryegrass
- RS = Ryegrass and straw