


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/C/7 Intensive Barley Growing - Cereals and Beans

Rothamsted Research

Rothamsted Research (1963) *62/R/C/7 Intensive Barley Growing - Cereals and Beans* ; Yields Of The Field Experiments 1962, pp 113 - 114 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/C/7.1

INTENSIVE BARLEY GROWING EXPERIMENT

Little Knott I - 1962, the second year

For treatments etc., see "The Numerical Results of the Field Experiments" 61/C/8.

Area of each plot (acres): 0.0212. Area harvested: 0.0140.

Cultivations, etc.: Ploughed: Sept 16, 1961.

Spring beans: Seed placement drilled at 200 lb per acre:
Feb 21, 1962. Combine harvested: Sept 21. Variety: Tick 30B.

Oats: 'Nitro-Chalk' applied: Feb 23, 1962. Seed combine drilled at 4 bushels per acre: Mar 2. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 24. Combine harvested: Aug 25. Variety: Condor.

Spring wheat: 'Nitro-Chalk' applied: Feb 23, 1962. Seed combine drilled at 3 bushels per acre: Mar 2. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 24. Combine harvested: Sept 13. Variety: Jufy I.

Barley: 'Nitro-Chalk' applied, seed combine drilled at $2\frac{1}{2}$ bushels per acre: Feb 22, 1962. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: May 24. Combine harvested: Aug 27. Variety: Proctor.

Winter wheat: Seed combine drilled at $2\frac{1}{2}$ bushels per acre: Oct 11, 1961. 'Nitro-Chalk' applied: Feb 23, 1962. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: Apr 24. Combine harvested: Sept 3. Variety: Cappelle.

Note: (1) One plot (non-continuous spring wheat preceding beans), which should have received 0.45 cwt N per acre, received 0.9 by mistake.
(2) Yields were only taken for sequences 3, 6, 7, 8, 9 and 10

Standard errors per plot. Grain (at 85% dry matter):

Winter wheat (9 and 10): 4.53 cwt per acre or 11.9% (7 d.f.)

Spring wheat (3, 6 and 8): 2.44 cwt per acre or 8.7% (14 d.f.)

62/C/7.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Winter wheat (9 and 10)

N: cwt per acre

Crop in 1961	None	0.3	0.6	0.9	Mean
	(± 3.20)				(± 1.60)
Spring wheat	23.9	24.2	35.6	39.2	30.7
Beans	39.0	45.5	46.1	51.3	45.5
Mean (± 2.26)	31.4	34.9	40.8	45.2	38.0

Mean dry matter % as harvested: 83.6

Spring wheat (3, 6, 8)

Previous crop	Spring wheat					Oats	Mean
	None	0.3	0.45* ⁺	0.6	0.9	0.45*	
N: cwt per acre	19.3	21.7	26.2	29.7	26.6	33.2	27.9
	(± 1.72)		(± 0.86)	(± 1.72)		(± 0.86)	

Mean dry matter % as harvested: 73.8

Barley 7

N: cwt per acre

None	0.3	0.6	0.9	Mean
33.0	42.9	42.9	46.8	41.4

Mean dry matter % as harvested: 82.6

*mean of 8 - others mean of 2

⁺includes 1 estimated value