


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1962

[Full Table of Content](#)



---

## 62/R/C/11 Levels of N and K - Cocksfoot 5th Year

### Rothamsted Research

Rothamsted Research (1963) *62/R/C/11 Levels of N and K - Cocksfoot 5th Year* ; Yields Of The Field Experiments 1962, pp 122 - 123 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/C/11.1

### GRASS

Levels of N and K - Harwoods Piece 1962 - the 5th year.

Design: 4 randomised blocks of 12 plots each.

Area of each plot: 0.0087 acres. Area harvested: 0.0059 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre as 'Nitro-Chalk'.

Potash: None; 0.3; 0.6 cwt  $K_2O$  per acre as muriate of potash.

All the above in the presence of 0.6 cwt  $P_2O_5$  per acre as superphosphate.

In addition 2 plots per block, receiving 0.9 cwt N and 0.6 cwt  $K_2O$  per acre, also received phosphate at either None or 1.2 cwt  $P_2O_5$  per acre as superphosphate.

Note: (1) N and K dressings are applied for each cut. All P dressings are applied once annually.

(2) All treatments were applied to the same plots as in the previous seasons.

Basal dressing: None.

Cultivations, etc.: Ryegrass sprayed with dalapon at 8 lb in 40 gallons per acre: Sept 7, 1961 and again at 4 lb in 40 gallons per acre: Sept 27. Ploughed: Nov 8. Ground chalk at 3 tons per acre applied to plots 1 - 24: Feb 21, 1962. Rotary cultivated: Apr 13. 1st dressing of fertilisers applied: Apr 16. Seed drilled at 40 lb per acre: Apr 24. Sprayed with MCPB/MCPA at 5 pints in 40 gallons per acre: June 5. Cut twice: Aug 13 and Oct 3. Variety: S37 Cocksfoot.

Note: (3) For details of the previous years' results see "Results of the Field Experiments" 58/Cg/2, 59/Cg/2, 60/Ci/1 and 61/Dg/1.

Standard errors per plot. Dry matter:

1st cut: 1.44 cwt per acre or 8.7% (33 d.f.)

2nd cut: 2.51 cwt per acre or 9.5% (33 d.f.)

Total of 2 cuts: 3.34 cwt per acre or 7.8% (33 d.f.)

62/C/11.2

Summary of Results

Dry matter: cwt per acre

cwt per acre														
N*	0.0	0.3	0.3	0.3	0.6	0.6	0.6	0.9	0.9	0.9	0.9	0.9		
P <sub>2</sub> O <sub>5</sub>	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.0	1.2	
K <sub>2</sub> O*	0.0	0.0	0.3	0.6	0.0	0.3	0.6	0.0	0.3	0.6	0.6	0.6	0.6	Mean
1st cut (±0.71)	8.5	14.1	17.5	18.0	14.2	18.3	20.1	13.5	18.0	19.8	17.6	19.4	16.6	
2nd cut (±1.25)	7.0	22.9	23.8	22.2	28.2	30.4	29.3	29.9	30.4	30.2	31.1	31.1	26.4	
Total of 2 cuts (±1.66)	15.5	37.1	41.2	40.2	42.4	48.7	49.3	43.4	48.4	50.0	48.8	50.6	42.9	

\*For each cut.

Mean dry matter % as cut:

1st cut: 18.3

2nd cut: 14.5

Total of 2 cuts: 16.4