


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1962

[Full Table of Content](#)



### 62/R/DA/4 Spring Wheat - Azotobacter Inoculation and N

#### Rothamsted Research

Rothamsted Research (1963) *62/R/DA/4 Spring Wheat - Azotobacter Inoculation and N* ; Yields Of The Field Experiments 1962, pp 128 - 128 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/Da/4

SPRING WHEAT

Effects of nitrogen and inoculation with Azotobacter - Fosters Corner 1962.

Design: 4 randomised blocks of 6 plots each.

Area of each plot: 0.0112 acres.

Treatments. All combinations of:-

Nitrogen: None; 0.2; 0.8 cwt N per acre applied as 'Nitro-Chalk'.

Azotobacter inoculation: None (sterile medium, no carbon source);

Azotobacter culture applied to seed.

Basal dressing: 2½ cwt compound fertiliser (20% P<sub>2</sub>O<sub>5</sub>, 20% K<sub>2</sub>O) per acre combine drilled.

Cultivations, etc.: Ploughed: Dec 16, 1961 - Jan 30, 1962. 'Nitro-Chalk' applied, seed drilled at 2¼ bushels per acre: Mar 17.

Sprayed with MCPA/MBA at 4 pints in 40 gallons per acre: June 2.

Combine harvested: Sept 13. Variety: Jufy I. Previous crops:

Spring wheat 1960; sugar beet 1961.

Note: Crop samples were taken during the season for counts of Azotobacter. Measurements of the height of the crop were made in July.

Standard error per plot.

Grain (at 85% dry matter): 2.96 cwt per acre or 7.6% (15 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Inoculation	N cwt per acre			Mean
	None	0.2	0.8	
		(±1.48)		(±0.85)
None	37.9	37.6	40.7	38.7
<u>Azotobacter</u>	39.3	39.3	40.2	39.6
Mean (±1.04)	38.6	38.4	40.4	39.1

Mean dry matter % as harvested: 72.4