


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/DC/1 Winter Beans - Residual Effects of Chalk

Rothamsted Research

Rothamsted Research (1963) *62/R/DC/1 Winter Beans - Residual Effects of Chalk* ; Yields Of The Field Experiments 1962, pp 130 - 130 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/Dc/1

WINTER BEANS

The residual effect of levels of chalk - Great Field I 1962, the 2nd year.

Design: 4 randomised blocks of 5 plots each.

Area of each plot: 0.0193 acres. Area harvested: 0.0117 acres.

Treatments: Ground chalk tons per acre applied in two dressings, half before ploughing on Dec 6, 1960 and half after ploughing on Dec 13, 1960:- None; 1; 2; 3; 4.

Basal dressing: $3\frac{1}{4}$ cwt per acre compound fertiliser (14% P_2O_5 , 28% K_2O) placement drilled.

Cultivations, etc.: Ploughed: Sept 18, 1961. Seed placement drilled at 275 lb per acre: Oct 10. Combine harvested: Sept 5, 1962. Variety: Garton's SQ. Previous crops: Winter wheat 1960, spring beans 1961.

Notes: (1) Samples were taken for counts of pods and beans.
(2) For the previous year's results see 'Numerical Results of the Field Experiments' 61/Dd/1.

Standard error per plot.

Grain (at 85% dry matter): 1.31 cwt per acre or 4.5% (12 d.f.)

Erratum to the 'Numerical Results of the Field Experiments' 1961 page 61/Dd/1. Area of each plot should read 0.0193 not 0.0212 acres.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Ground chalk: tons per acre applied Dec, 1960					Mean
None	1	2	3	4	
26.3	28.1	29.9	31.9	30.7	29.4
(±0.65)					

Mean dry matter % as harvested: 76.8