


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1962

[Full Table of Content](#)



---

## 62/R/DD/3 Potatoes - Fungicides

### Rothamsted Research

Rothamsted Research (1963) *62/R/DD/3 Potatoes - Fungicides* ; Yields Of The Field Experiments 1962, pp 139 - 140 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/Dd/3.1

## POTATOES

Control of blight (*Phytophthora infestans*) by copper and tin fungicides - Long Hoos V 1962.

Design: Two 6 x 6 Latin squares, one for each variety, - King Edward and Ulster Supreme.

Area of each plot: 0.0129 acres. Area harvested: 0.0077 acres.

### Treatments:

No fungicide (0)

Sprayed twice with:-

Commercial copper oxychloride wettable powder at  $2\frac{1}{2}$  lb Cu per acre (1)

Copper oxychloride at  $2\frac{1}{2}$  lb Cu per acre with either 0.25 (2), 0.50 (3), or 1.00% wax (4)

Commercial triphenyltin acetate wettable powder at 0.75 lb triphenyltin acetate per acre (5).

All sprays applied in 100 gallons per acre.

Basal dressing: 10 cwt compound fertiliser (10% N, 10%  $P_2O_5$ , 18%  $K_2O$ ) per acre.

Cultivations, etc.: Sprayed with dalapon at 8 lb in 40 gallons per acre: Sept 7, 1961 and again at 4 lb in 40 gallons per acre: Sept 26. Ploughed: Oct 17. Rotary cultivated, basal dressing applied, potatoes machine planted: Apr 24, 1962. Earthed up: July 7. Copper and tin fungicides applied: King Edward - July 20 and Aug 9, Ulster Supreme - July 25 and Aug 22. King Edward lifted: Oct 11. Ulster Supreme haulm destroyed mechanically, crop lifted: Oct 15. Previous crops: Spring beans 1960; spring wheat 1961.

Note: The destruction of foliage by blight and the percentage weight of infected tubers at harvest were estimated.

Standard errors per plot. Total tubers tons per acre:

King Edward: 0.626 tons per acre or 4.8% (20 d.f.)

Ulster Supreme: 1.027 tons per acre or 7.5% (20 d.f.)



62/Da/3.2

Summary of Results

Variety	Fungicide						Mean
	0	1	2	3	4	5	
<u>Total tubers: tons per acre</u>							
<u>King Edward</u>							
Mean ( $\pm 0.256$ )	12.43	12.66	13.04	13.21	12.89	14.04	13.05
Increase ( $\pm 0.361$ )		0.23	0.61	0.78	0.46	1.61	
<u>Ulster Supreme</u>							
Mean ( $\pm 0.419$ )	12.02	13.20	13.57	13.79	13.95	15.45	13.67
Increase ( $\pm 0.593$ )		1.18	1.55	1.77	1.93	3.43	
<u>Percentage ware (1½" riddle)</u>							
<u>King Edward</u>							
Mean	88.3	89.5	90.0	88.9	89.9	90.3	89.5
Increase		1.2	1.7	0.6	1.6	2.0	
<u>Ulster Supreme</u>							
Mean	97.7	97.9	97.9	97.7	97.8	98.1	97.8
Increase		0.2	0.2	0.0	0.1	0.4	

Fungicide.

- 0 = No fungicide
- 1 = Commercial copper oxychloride wettable powder at 2½ lb Cu per acre.
- 2 = Copper oxychloride at 2½ lb Cu per acre with 0.25% wax.
- 3 = " " " " " " " " " 0.50% wax.
- 4 = " " " " " " " " " 1.00% wax.
- 5 = Commercial triphenyltin acetate wettable powder at 0.75 lb triphenyltin acetate per acre.