


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/A/1 Broadbalk - Wheat

Rothamsted Research

Rothamsted Research (1963) *62/R/A/1 Broadbalk - Wheat* ; Yields Of The Field Experiments 1962, pp 5 - 6 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/A/1.1

WHEAT - BROADBALK 1962

The 119th year

For history, treatments, etc., see "Details of the Classical and Long Term Experiments" 1956.

Cultivations, etc.:

Cropped sections. Dung and ground chalk applied, all plots ploughed: Sept 7, 1961. Autumn fertilisers applied: Sept 29. Seed drilled at 3 bushels per acre: Nov 1. Spring fertilisers applied: Apr 17, 1962. Section IA sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: Apr 24. Second dressing of nitrate of soda applied to plot 16: May 4. Combine harvested: Sept 15. Variety: Squarehead's Master 13/4.

Fallow section. (II) Ploughed: Sept 7, 1961; May 16, 1962; July 16.

Broadbalk Wilderness. N.

Cultivations, etc.: Shrubs grubbed out: Nov 17 - 30, 1961. Part grazed: Dec 12 - 13; May 11 - 30, 1962; June 12 - 18; July 27 - Aug 1; Sept 6 - 26. Grass topped: May 23; Aug 1; Sept 26.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Section Years after fallow	IB	III	IV	VA	VB	IA	Mean
	1	2	3	unlimed 4	limed 4	11	
2A	40.9	22.2	26.0	24.3	23.3	26.9	26.5
2B	37.5	25.9	26.7	25.2	24.2	27.0	27.4
3	18.5	11.7	9.5	11.2	10.8	12.5	11.9
5	23.1	11.8	12.4	12.4	11.4	13.1	13.6
6	23.0	16.0	14.8	12.2	14.1	17.3	15.9
7	29.4	23.1	21.4	16.6	15.3	22.7	21.5
8	26.5	32.4	28.1	24.6	20.6	31.9	27.7
9	23.8	19.3	17.5	14.6	12.9	22.6	18.1
10	19.7	24.2	21.8	20.6	18.1	20.9	21.4
11	22.9	24.9	20.3	19.0	15.8	28.2	21.6
12	22.2	24.6	19.0	19.5	17.3	25.3	21.2
13	27.6	20.7	17.1	17.1	15.7	26.4	19.9
14	25.0	24.9	19.7	19.2	16.9	26.7	21.8
15	24.0	20.2	19.0	16.8	12.0	19.6	18.8
16	24.2	27.7	25.8	21.9	21.7	27.2	25.1
17	24.8	25.1	21.3	19.8	18.4	24.8	22.4
18	17.2	10.7	8.9	9.4	9.9	12.9	11.0
19	25.0	20.4	15.0	13.0	14.4	21.9	17.9
20	16.9					14.9	16.2

Mean dry matter % as harvested: 78.6

62/A/1.2

Straw (at 85% dry matter): cwt per acre

Section Years after fallow	IB	III	IV	VA	VB	IA	Mean
	1	2	3	unlimed	limed	11	
				4	4		
2A	45.5	32.8	29.8	24.9	24.8	30.5	31.4
2B	44.4	38.6	32.7	32.9	26.5	26.9	34.6
3	15.8	13.0	8.2	10.6	9.3	10.7	11.1
5	27.7	13.6	12.4	15.6	14.1	12.7	15.5
6	27.5	18.8	17.0	15.9	18.1	20.6	19.1
7	36.5	27.0	30.3	28.6	25.7	25.3	29.1
8	35.8	38.4	34.5	33.0	28.4	36.8	34.8
9	27.7	27.8	27.4	22.0	21.1	30.1	26.1
10	16.8	25.5	20.0	23.1	13.5	22.0	20.7
11	20.9	26.4	18.0	19.6	13.5	25.4	20.7
12	21.8	23.3	19.9	20.7	17.4	25.7	21.2
13	35.3	25.4	24.5	25.3	23.6	31.8	26.7
14	26.4	25.4	20.8	22.8	19.2	30.0	23.4
15	31.9	22.4	19.2	21.5	11.8	24.7	21.4
16	31.2	37.3	32.2	28.0	28.8	39.6	32.9
17	29.4	29.4	26.7	24.9	24.4	30.0	27.4
18	20.4	9.9	10.2	9.3	9.5	13.1	11.5
19	30.4	22.5	18.6	16.3	17.3	24.9	21.2
20	20.8					17.2	19.6

Mean dry matter % as harvested: 87.2