


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1962

[Full Table of Content](#)



62/R/B/9 N Levels and Residues Two Course Rotation

Rothamsted Research

Rothamsted Research (1963) *62/R/B/9 N Levels and Residues Two Course Rotation* ; Yields Of The Field Experiments 1962, pp 91 - 92 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/B/9.1

N LEVELS AND RESIDUES ROTATION

Direct and residual effects of sulphate of ammonia - Long Hoos III 1962, the 3rd year.

Rotation: Wheat, potatoes.

Design (each crop): 3 randomised blocks of 9 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments. All combinations of:-

Nitrogen (applied as sulphate of ammonia) at 3 levels in 1961.

Nitrogen at 3 levels in 1960, and repeated on the same plots in 1962.

To wheat: None; 0.5; 1.0 cwt N per acre.

To potatoes: None; 0.75; 1.50 cwt N per acre.

Basal dressing (per acre):

To wheat: $2\frac{1}{4}$ cwt compound fertiliser, 14% P_2O_5 , 28% K_2O combine drilled.

To potatoes: 5 cwt compound fertiliser, 14% P_2O_5 , 28% K_2O broadcast on the flat.

Cultivations, etc.:

Wheat: Ploughed: Oct 6, 1961. Rotary cultivated: Oct 9. Seed drilled at $2\frac{1}{2}$ bushels per acre: Oct 12. Sulphate of ammonia applied: Apr 9, 1962. Sprayed with MCPA/TBA at 4 pints in 40 gallons per acre: Apr 24. Combine harvested: Sept 3. Variety: Cappelle.

Potatoes: Ploughed: Oct 12, 1961. Rotary cultivated: Mar 26, 1962. Basal fertiliser and sulphate of ammonia applied: Apr 11. Ridged, potatoes planted: Apr 13. Earthed up: July 7. Sprayed with maneb at $1\frac{1}{2}$ lb in 18 gallons per acre: July 19. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 20 gallons per acre: Aug 10 and again Aug 24. Sprayed with undiluted BOV at 15 gallons per acre: Sept 25. Lifted: Oct 11. Variety: Ulster Supreme.

Standard errors per plot.

Wheat. Grain (at 85% dry matter): 2.59 cwt per acre or 6.7% (16 d.f.)

Potatoes. Total tubers: 0.790 tons per acre or 9.4% (16 d.f.)

Note: For details of the previous years results see "Results of the Field Experiments" 60/B/10 and 61/B/9.

62/B/9.2

Summary of Results

Wheat, Grain (at 85% dry matter): cwt per acre

	N: cwt per acre in 1960 & 1962			Mean
	None	0.5	1.0	
N: cwt per acre to potatoes in 1961		(±1.49)		(±0.86)
None	28.2	34.2	42.2	34.8
0.75	30.6	40.2	44.4	38.4
1.50	37.1	43.9	48.4	43.1
Mean (±0.86)	32.0	39.4	45.0	38.7

Mean dry matter % as harvested: 84.0

Potatoes

	N: cwt per acre in 1960 & 1962			Mean
	None	0.75	1.50	
N: cwt per acre to wheat in 1961	<u>Total tubers: tons per acre</u>			
		(±0.455)		(±0.263)
None	4.95	8.95	11.64	8.52
0.5	4.60	8.90	10.76	8.08
1.0	4.94	9.42	11.53	8.63
Mean (±0.263)	4.83	9.09	11.31	8.40

	N: cwt per acre to wheat in 1961			
	<u>Percentage ware (1½" riddle)</u>			
None	94.6	96.6	96.7	95.9
0.5	94.6	96.9	97.6	96.4
1.0	94.9	96.1	97.2	96.1
Mean	94.7	96.5	97.2	96.1