


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1962

[Full Table of Content](#)



---

## 62/R/B/10 Weedkiller-cultivation Rotation

### Rothamsted Research

Rothamsted Research (1963) *62/R/B/10 Weedkiller-cultivation Rotation* ; Yields Of The Field Experiments 1962, pp 93 - 96 - DOI: <https://doi.org/10.23637/ERADOC-1-164>

62/B/10.1

### WEEDKILLER CULTIVATION ROTATION

Great Harpenden I 1962, the second year

A comparison of weed control by various cultivation methods and by pre-emergence weedkillers.

For previous history, rotation etc., see "Numerical Results of the Field Experiments" 61/B/10.

Area harvested (acres): Beans - 0.0201; Wheat, potatoes, barley - 0.0107 (sub plots).

Prometryne, at  $2\frac{1}{2}$  lb active material in 20 gallons per acre, has replaced simazine as the weedkiller on potatoes.

Basal dressings per acre:- Spring wheat: 3 cwt compound fertiliser (16% N, 9%  $P_2O_5$ , 9%  $K_2O$ ) combine drilled. All other crops - as 1961.

#### Operations in 1962

Note: Spring beans and spring wheat were again sown instead of winter beans and winter wheat.

Cultivations, etc.: All plots except barley plots sprayed with dalapon at 8 lb in 40 gallons per acre: Sept 7, 1961 and again at 4 lb in 40 gallons per acre: Sept 26.

Spring beans: T plots rigid-tine cultivated 3 times: Nov 1, 3 and 4, 1961. P and reserve plots ploughed: Nov 6. R plots rotary cultivated: Nov 23. All plots except R plots spring-tine cultivated twice: Feb 23 and Mar 13, 1962. R plots rotary cultivated: Mar 13. Seed drilled at 200 lb per acre, all plots harrowed: Mar 14. X plots sprayed with simazine: Apr 11. M and reserve plots mechanically weeded: May 2. M and reserve plots tractor hoed 3 times: May 10, June 1 and June 20. Combine harvested: Sept 20. Variety: Tick 30B.

Spring wheat: T plots rigid-tine cultivated 3 times: Nov 1, 3 and 4, 1961. P and reserve plots ploughed: Nov 6. R plots rotary cultivated: Nov 23. Spring-tine cultivated twice (except R plots): Mar 2 and 15, 1962. R plots rotary cultivated, seed drilled at 3 bushels per acre, all plots harrowed: Mar 16. Rolled: Apr 30. H sub plots and reserve plots sprayed with MCPA/MBA at 4 pints in 40 gallons per acre: May 28. Combine harvested: Sept 14. Variety: Jufy I.

Potatoes: T plots rigid-tine cultivated 3 times: Nov 1, 3 and 4, 1961. P and reserve plots ploughed: Nov 6. R plots rotary cultivated: Nov 22. Spring-tine cultivated twice (except R plots), R plots rotary cultivated once: Mar 22, 1962. Basal compound fertiliser applied: Mar 23. Seed machine planted: Mar 27. Rolled: Mar 28. Mechanically weeded (except X plots): May 1. Chain harrowed (except X plots): May 2.



62/B/10.2

Grubbed (except X plots): May 4. X plots sprayed with prometryne: May 23. Reridged (except X plots): May 31. Y plots sprayed with prometryne: June 2. Mechanically weeded (except X and Y plots): June 7. M and reserve plots grubbed: June 18. M, reserve plots and E sub plots of X plots grubbed and ridged: July 5. Sprayed with maneb at  $1\frac{1}{2}$  lb active material in 18 gallons per acre: July 20. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 20 gallons per acre: Aug 10. Sprayed with undiluted BOV at 15 gallons per acre: Sept 25. Lifted: Oct 12. Variety: Majestic.

Barley: T plots rigid-tine cultivated twice, P and reserve plots ploughed: Oct 17, 1961. R plots rotary cultivated: Nov 22. Spring-tine cultivated twice (except R plots): Feb 23 and Mar 15, 1962. R plots rotary cultivated, seed drilled at  $2\frac{1}{4}$  bushels per acre, all plots harrowed: Mar 16. Rolled: Apr 30. H sub plots and reserve plots sprayed with MCPA/MBA at 4 pints in 40 gallons per acre: May 28. Combine harvested: Aug 27. Variety: Proctor.

Standard errors per plot.

Spring beans, grain (at 85% dry matter): 3.84 cwt per acre or 13.2% (15 d.f.)

Spring wheat, grain (at 85% dry matter)

Whole plot: 2.49 cwt per acre or 7.6% (15 d.f.)

Sub plot: 1.27 cwt per acre or 3.9% (18 d.f.)

Potatoes, total tubers

Whole plot: 1.458 tons per acre or 10.3% (12 d.f.)

Sub plot: 0.900 tons per acre or 6.3% (15 d.f.)

Barley, grain (at 85% dry matter)

Whole plot: 3.21 cwt per acre or 7.8% (12 d.f.)

Sub plot: 1.60 cwt per acre or 3.9% (15 d.f.)

### Summary of Results

Beans, grain (at 85% dry matter): cwt per acre

Treatment after planting	Initial cultivation			Mean
	P	R	T	
M ( $\pm 2.72$ )	27.0	31.1	27.8	28.6 ( $\pm 1.57$ )
X ( $\pm 1.92$ )	29.4	30.0	28.1	29.2 ( $\pm 1.11$ )
Mean ( $\pm 1.57$ )	28.6	30.3	28.0	29.0

Reserve plots 29.3 ( $\pm 1.57$ )

General mean: 29.1

Mean dry matter % as harvested: 78.0



62/B/10.3

Spring wheat, grain (at 85% dry matter): cwt per acre

	Initial cultivation			Mean
	P	R	T	
Mean ( $\pm 1.02$ )	31.8	33.7	34.0	33.2
<u>Treatment in 1961</u>				
M ( $\pm 1.76$ )	32.5	33.2	35.1	33.6 ( $\pm 1.02$ )
X ( $\pm 1.24$ )	31.5	34.0	33.4	33.0 ( $\pm 0.72$ )
<u>Spray in 1962</u>				
- ( $\pm 1.08$ ) <sup>(2)</sup>	33.2	33.7	35.4	34.1
H	30.5	33.8	32.5	32.3
Diff ( $\pm 0.74$ )	-2.7	+0.1	-2.9	-1.8 ( $\pm 0.42$ )

Reserve plots: 30.8 ( $\pm 1.02$ )

General mean: 32.6

Mean dry matter % as harvested: 74.2

Potatoes

Treatment after planting	Initial cultivation				Mean	Not earthed up	Earthed up
	P	R	T	Mean			
<u>Total tubers: tons per acre</u>							
	( $\pm 1.031$ )			( $\pm 0.595$ )	( $\pm 0.367$ ) <sup>(1)</sup> ( $\pm 0.649$ ) <sup>(2)</sup>		
M	14.97	15.44	14.44	14.95	14.92	14.98	
X	14.09	12.63	12.15	12.95	13.17	12.74	
Y	13.11	15.51	14.29	14.30			
Mean ( $\pm 0.595$ )	14.06	14.53	13.62	14.07			

Reserve plots: 14.56 ( $\pm 0.595$ )

General mean: 14.19

(1) For use in vertical and interaction comparisons

(2) For use in horizontal and diagonal comparisons



62/B/10.4

Potatoes

Treatment after planting	Initial cultivation				Not earthed up	Earthed up
	P	R	T	Mean		
	<u>Percentage ware (1½" riddle)</u>					
M	96.4	97.1	96.4	96.6	96.2	97.1
X	96.9	97.2	96.8	97.0	97.1	96.8
Y	95.7	96.3	95.7	95.9		
Mean	96.3	96.9	96.3	96.5		

Reserve plots: 95.8

General mean: 96.3

Barley, grain (at 85% dry matter): cwt per acre

	Initial cultivation			
	P	R	T	Mean
Mean ( $\pm 1.31$ )	41.8	39.8	38.3	40.0
<u>Treatment in 1961</u>		( $\pm 2.27$ )		( $\pm 1.31$ )
M	40.5	41.0	38.3	39.9
X	44.3	40.3	36.9	40.5
Y	40.6	38.3	39.8	39.6
<u>Spray in 1962</u>				
- ( $\pm 1.39$ )*	42.8	41.0	38.8	40.8
H	40.8	38.7	37.9	39.1
Diff ( $\pm 0.92$ )	-2.0	-2.3	-0.9	-1.7 ( $\pm 0.53$ )

Reserve plots: 43.9 ( $\pm 1.31$ )

General mean: 41.0

Mean dry matter % as harvested: 82.2

\* For use in horizontal and diagonal comparisons.