


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1963

[Full Table of Content](#)



---

## 63/R/CW/B/10 Weedkiller-cultivation Rotation

### Rothamsted Research

Rothamsted Research (1964) *63/R/CW/B/10 Weedkiller-cultivation Rotation* ; Yields Of The Field Experiments 1963, pp 117 - 121 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/B/10.1

CULTIVATION - WEEDKILLER ROTATION

(CW)

Great Harpenden I 1963, the third year

A comparison of weed control by various cultivation methods and by pre-emergence weedkillers.

For previous history, rotation treatments etc., see 'Numerical Results of the Field Experiments' 61/B/10 and 62/B/10.

Area harvested (acres): Winter beans - 0.0201. Spring wheat, potatoes, barley - 0.0107.

The weedkiller used on potatoes is now a mixture of 2 lb prometryne and 0.75 lb paraquat in 40 gallons per acre. X and Y treatments for potatoes were identical.

Basal dressings per acre: Spring wheat and barley: 2.5 cwt compound fertiliser, 20% N, 10%  $P_{2}O_{5}$ , 10%  $K_{2}O$ . Potatoes and beans, as 1961.

Operations in 1963

Note: Spring wheat was again sown instead of winter wheat.

Cultivations, etc.:

Winter beans: R plots rotary cultivated, T plots rigid-tine cultivated 3 times: Oct 22, 1962. R plots rotary cultivated second time: Oct 23. P and reserve plots ploughed: Oct 25. T, P and reserve plots spring-tine cultivated twice: Oct 29, Nov 30. Seed drilled at 275 lb per acre, all plots spring-tine cultivated: Dec 1. X and Y\* plots sprayed with simazine: Dec 19. Y\* plots sprayed with simazine: Apr 19, 1963. M and reserve plots tractor hoed twice: May 21, May 29. All plots sprayed with menazon at 5.6 fluid oz in 40 gallons per acre: June 22. Combine harvested: Oct 5. Variety: Pedigree.

Spring wheat: Sprayed with dalapon at 7.4 lb in 40 gallons per acre: Oct 20, 1962 and again at 3.7 lb in 40 gallons per acre: Oct 30. T plots rigid-tine cultivated twice, P and reserve plots ploughed: Apr 3, 1963. T, P and reserve plots spring-tine cultivated: Apr 18. R plots rotary cultivated, T, P and reserve plots spring-tine harrowed, seed drilled at 3 bushels per acre: Apr 19. Rolled: May 8. H sub plots and reserve plots sprayed with methoxychloro-benzoic acid/MCPA (MBA/MCPA) at 4 pints in 40 gallons per acre: June 6. Combine harvested: Sept 19. Variety: Jufy I.

Potatoes: R plots rotary cultivated, T plots rigid-tine cultivated twice: Oct 22, 1962. P and reserve plots ploughed: Oct 25. T, P and reserve plots spring-tine cultivated: Apr 25, 1963. R plots

\* half-rate

63/B/10.2

rotary cultivated: Apr 27. Basal compound fertiliser applied: Apr 30. T, P and reserve plots spring-tine cultivated, seed machine planted: May 3. Rolled: May 6. X and Y plots sprayed with prometryne at 2 lb plus paraquat at 0.75 lb in 40 gallons per acre, M plots chain harrowed: May 31. M and reserve plots grubbed: June 7. M and reserve plots mechanically weeded and grubbed: June 17. Reserve plots earthed up: June 26. M and ME sub-plots earthed up (round and pointed ridges), E sub-plots of X and Y plots grubbed twice and earthed up: June 28. Sprayed with maneb at 1.2 lb in 20 gallons per acre: July 19. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 20 gallons per acre: Sept 4. Sprayed with undiluted BOV at 16 gallons per acre: Sept 13. Lifted: Oct 23. Variety: Majestic.

Barley: All plots spring-tine cultivated twice: Nov 27, 1962, Mar 27, 1963. T plots rigid-tine cultivated: Apr 3. P and reserve plots ploughed: Apr 4. All plots except R plots disc-harrowed: Apr 11. R plots rotary cultivated, P and reserve plots spring-tine cultivated: Apr 19. T, P and reserve plots spring-tine harrowed, seed drilled at 2 bushels per acre: Apr 20. Rolled: May 8. H sub-plots and reserve plots sprayed with methoxychloro-benzoic acid/MCPA (MBA/MCPA) at 4 pints in 40 gallons per acre: June 6. Combine harvested: Sept 7. Variety: Proctor.

Standard errors per plot:

Winter beans, grain (at 85% dry matter): 3.63 cwt per acre or 10.9%  
(9 d.f.)

Spring wheat, grain (at 85% dry matter):

Whole plot: 2.42 cwt per acre or 6.6% (15 d.f.)

Sub plot: 2.00 cwt per are or 5.4% (18 d.f.)

Potatoes, total tubers:

Whole plot: 0.752 tons per acre or 5.5% (15 d.f.)

Sub plot: 0.456 tons per acre or 3.3% (18 d.f.)

Barley, grain (at 85% dry matter):

Whole plot: 1.91 cwt per acre or 4.5% (12 d.f.)

Sub plot: 3.28 cwt per acre or 7.8% (15 d.f.)



63/B/10.3

Summary of Results

Beans, grain (at 85% dry matter): cwt per acre

Treatment after planting	Initial cultivation			Mean
	P	R	T	
		(±2.57)		(±1.48)
M	39.1	27.5	31.5	32.7
X	33.8	33.9	29.7	32.4
Y	33.0	32.5	34.8	33.4
Mean (±1.48)	35.3	31.3	32.0	32.8

Reserve plots: 36.5 (±1.82)

General mean: 33.5

Mean dry matter % as harvested: 68.2

Spring wheat, grain (at 85% dry matter): cwt per acre

	Initial cultivation			Mean
	P	R	T	
Mean (±0.99)	36.3	36.7	35.6	36.2
<u>Treatment in 1962</u>				
M (±1.71)	35.4	36.4	36.4	36.1 (±0.99)
X (±1.21)	36.8	36.8	35.2	36.2 (±0.70)
<u>Spray in 1963</u>		(±1.14)*		
-	37.5	36.3	35.4	36.4
H	35.2	37.1	35.7	36.0
Diff (±1.15)	-2.3	+0.8	+0.3	-0.4 (±0.67)

Reserve plots: 38.8 (±0.99)

General mean: 36.8

Mean dry matter % as harvested: 79.0

\* For use in horizontal and diagonal comparisons only

63/B/10.4

Potatoes

Treatment after planting	Initial cultivation			Mean	Not earthed up	Earthed up
	P	R	T			
<u>Total tubers: tons per acre</u>						
					(±0.186)(1)	(±0.293)(2)
M (±0.532)	14.58	13.37	13.88	13.94 (±0.307)	13.90	13.98
X (±0.376)	13.40	13.90	13.84	13.70 (±0.217)	13.48	13.94
					(±0.132)(1)	(±0.262)(2)
Mean (±0.307)	13.79	13.72	13.85	13.79		
Reserve plots:	13.77 (±0.307)					
General mean:	13.78					

Percentage ware (1.5 inch riddle)

M	94.8	95.6	95.8	95.4	95.1	95.7
X	95.0	94.8	95.2	95.0	94.7	95.2
Mean	94.9	95.1	95.4	95.1		

Reserve plots: 95.9  
General mean: 95.3

- (1) For use in vertical and interaction comparisons  
(2) For use in horizontal and diagonal comparisons

63/B/10.5

Barley, grain (at 85% dry matter): cwt per acre

	Initial cultivation			Mean
	P	R	T	
Mean ( $\pm 0.78$ )	42.9	39.8	41.9	41.5
<u>Treatment in 1962</u>		( $\pm 1.35$ )		( $\pm 0.78$ )
M	42.8	39.2	43.6	41.8
X	41.4	40.8	41.6	41.2
Y	44.6	39.4	40.5	41.5
<u>Spray in 1963</u>		( $\pm 1.22$ )*		
-	42.8	40.7	40.2	41.2
H	43.0	38.8	43.6	41.8
Diff ( $\pm 1.89$ )	+0.2	-1.9	+3.4	+0.6 ( $\pm 1.09$ )

Reserve plots: 44.0 ( $\pm 0.78$ )

General mean: 42.1

Mean dry matter % as harvested: 71.8

\* For use in horizontal and diagonal comparisons