


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/LM/C/1 and 63/W/WAC/C/1 Effect of K and Mg - Clover

Rothamsted Research

Rothamsted Research (1964) *63/R/LM/C/1 and 63/W/WAC/C/1 Effect of K and Mg - Clover ; Yields Of The Field Experiments 1963*, pp 127 - 132 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/C/1.1

EFFECT OF K AND Mg

(LM and WAC)

K and Mg - Rothamsted (R) Sawyers I 1963 the fifth year and Woburn (W) Stackyard Series C 1963 the fourth year, - clover (sown in 1962 on both experiments).

Design: Sawyers I (R): 8 randomised blocks of 9 plots each.
Stackyard Series C (W): 4 randomised blocks of 9 plots each.

Area of each plot (acres):	Area harvested (acres):
Sawyers I (R): 0.0209	0.0045
Stackyard Series C (W): 0.0011	0.0003 - 0.0005

Treatments. All combinations of:-

Mg: None, 29, 58 lb Mg per acre applied as magnesium sulphate on
Sawyers I (R) and as kieserite on Stackyard Series C (W).

K: Sawyers I (R): 24, 95, 165 lb K per acre in 1962.
None, 71, 142 lb K per acre in 1963.

Stackyard Series C (W): None, 95, 190 lb K per acre.

All K as sulphate of potash, applied in 1962 and 1963.

In addition in 1962 magnesium-free calcium carbonate was applied to
blocks on Sawyers I (R) as in 1959, at 38, 76 cwt per acre.

Basal dressings per acre:

Sawyers I (R): 1.0 cwt P₂O₅ as triple superphosphate applied
in seedbed 1962, None in 1963.

Stackyard Series C (W): 1.0 cwt P₂O₅ as triple superphosphate in
spring 1963.

Cultivations, etc.:

Sawyers I (R): Magnesium-free calcium carbonate applied: At 30, 60
cwt per acre - Nov 21, 1961, at 8, 16 cwt per acre - Feb 19, 1962.
Magnesium sulphate and sulphate of potash applied: Mar 13, 1963.
Cut 3 times: June 11, July 29, Sept 26. Variety: Dorset Marl
Red Clover.

Stackyard Series C (W): Treatments and basal dressing applied:
Mar 21, 1963. Cut 3 times: June 13, July 25, Oct 11.
Variety: Dorset Marl Red Clover.

Note: For details of the previous years' results see 'Results of the
Field Experiments' 60/C1/3, 61/C/7 and 62/C/6.

63/C/1.2

Standard errors per plot. Clover, dry matter:

Sawyers I (R)

1st cut:	2.73 cwt per acre or 8.5% (48 d.f.)
2nd cut:	1.90 cwt per acre or 9.0% (48 d.f.)
3rd cut:	1.38 cwt per acre or 18.9% (48 d.f.)
Total of 3 cuts:	4.51 cwt per acre or 7.4% (48 d.f.)

Stackyard Series C (W)

1st cut:	2.98 cwt per acre or 10.8% (24 d.f.)
2nd cut:	1.33 cwt per acre or 9.1% (24 d.f.)
3rd cut:	0.91 cwt per acre or 9.1% (24 d.f.)
Total of 3 cuts:	3.98 cwt per acre or 7.6% (24 d.f.)

Summary of Results

Sawyers I (R)

Clover, Dry matter: cwt per acre

K: lb per acre 1962	24	95	165	Mg: lb per acre			
K: lb per acre 1963	None	71	142	None	29	58	Mean

<u>1st cut</u>							
Calcium carbonate cwt per acre	(±0.79)*			(±0.79)*			
38	27.4	34.2	36.3	32.8	33.1	32.0	32.6
76	24.5	34.0	37.3	31.0	32.9	31.9	31.9
Diff.	-2.9	-0.2	+1.0	-1.8	-0.2	-0.1	-0.7
	(±1.12)**			(±1.12)**			

<u>K: lb per acre</u>							
1962		1963		(±0.97)			
24	None	25.1	27.7	25.0	26.0		
95	71	33.8	34.5	34.0	34.1		
165	142	36.8	36.9	36.7	36.8		
Mean (±0.56)		31.9	33.0	31.9	32.3		

*For use in horizontal and interaction comparisons only.

** For use only in testing the differences of 2 differences.

Mean dry matter % as cut: 1st cut 19.4

63/c/1.3

Sawyers I (R)

Clover, Dry matter: cwt per acre

K: 1b per acre 1962	24	95	165	Mg: 1b per acre		
K: 1b per acre 1963	None	71	142	None	29	58

<u>2nd cut</u>						
Calcium carbonate cwt per acre	K: 1b per acre			(±0.55)*		
38	17.2	22.6	24.7	21.6	21.5	21.4
76	16.1	21.9	23.9	20.0	21.4	20.5
Diff.	-1.1	-0.7 (±0.78)**	-0.8	-1.6	-0.1 (±0.78)**	-0.9
K: 1b per acre						
	1962	1963		(±0.67)		(±0.39)
	24	None	16.2	17.2	16.5	16.6
	95	71	22.8	21.9	22.2	22.3
	165	142	23.6	25.4	24.1	24.3
Mean	(±0.39)		20.8	21.5	20.9	21.1

* For use in horizontal and interaction comparisons only.

** For use only in testing the difference of 2 differences.

Mean dry matter % as cut: 2nd cut 21.6

63/c/1.4

Sawyers I (R)

Clover, Dry matter: cwt per acre

K: lb per acre 1962	24	95	165	Mg: lb per acre			
K: lb per acre 1963	None	71	142	None	29	58	Mean

Calcium carbonate cwt per acre	<u>3rd cut</u>						Mean (±0.28)
	(±0.40)*			(±0.40)*			
38	6.6	7.9	8.2	7.8	7.7	7.2	7.5
76	5.0	7.4	8.5	6.7	6.9	7.4	7.0
Diff.	-1.6	-0.5 (±0.56)**	+0.3	-1.1	-0.8 (±0.56)**	+0.2	-0.5
K: lb per acre	1962	1963			(±0.49)		(±0.28)
	24	None	5.8	6.1	5.5	5.8	
	95	71	7.6	7.6	7.8	7.7	
	165	142	8.4	8.1	8.6	8.4	
Mean (±0.28)			7.2	7.3	7.3	7.3	

* For use in horizontal and interaction comparisons only.

** For use only in testing the difference of 2 differences.

Mean dry matter % as cut: 3rd cut 16.9

63/C/1.5

Sawyers I (R)

Clover, Dry matter: cwt per acre

K: lb per acre 1962	24	95	165	Mg: lb per acre		
K: lb per acre 1963	None	71	142	None	29	58

Calcium carbonate cwt per acre	Total of 3 cuts					
	1962		1963		Mean	
38	51.1	64.7	69.2	62.2	62.3	60.6
76	45.7	63.3	69.7	57.7	61.2	59.8
Diff.	-5.4	-1.4 (±1.84)**	+0.5	-4.5	-1.1 (±1.84)**	-0.8
						-2.1
K: lb per acre						
			(±1.59)		(±0.92)	
1962	24	None	47.1	51.1	47.1	48.4
	95	71	64.1	63.9	64.0	64.0
	165	142	68.7	70.3	69.5	69.5
Mean	(±0.92)		60.0	61.8	60.2	60.6

* For use in horizontal and interaction comparisons only.

** For use only in testing the difference of 2 differences.

Mean dry matter % as cut: Total of 3 cuts 19.3

63/C/1.6

Stackyard Series C (W)

Clover, Dry matter: cwt per acre

K: lb per acre in 1962 and 1963	Mg: lb per acre			Mean	Mg: lb per acre			Mean
	None	29	58		None	29	58	
<u>1st cut</u>								
		(±1.49)		(±0.85)		(±0.66)		(±0.38)
None	15.6	17.5	17.7	16.9	8.1	8.4	7.8	8.1
95	32.5	32.3	33.4	32.7	17.6	16.6	17.1	17.1
190	31.5	33.3	33.5	32.7	17.4	18.8	19.5	18.6
Mean	26.5	27.7	28.2	27.5	14.4	14.6	14.8	14.6
		(±0.85)			(±0.38)			
<u>3rd cut</u>								
		(±0.45)		(±0.26)		(±1.99)		(±1.15)
None	4.9	3.7	3.9	4.2	28.6	29.5	29.4	29.2
95	10.4	11.6	11.6	11.2	60.5	60.6	62.1	61.0
190	14.6	15.0	14.4	14.7	63.5	67.1	67.4	66.0
Mean	10.0	10.1	10.0	10.0	50.8	52.4	53.0	52.1
		(±0.26)			(±1.15)			

Mean dry matter % as cut:	1st cut	16.2
	2nd cut	16.8
	3rd cut	17.8
	Total of 3 cuts	16.9