


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1963

[Full Table of Content](#)



### 63/R/LL/C/3 and 63/W/WLL/C/3 Long-term Liming - Spring Beans

#### Rothamsted Research

Rothamsted Research (1964) *63/R/LL/C/3 and 63/W/WLL/C/3 Long-term Liming - Spring Beans* ; Yields Of The Field Experiments 1963, pp 135 - 138 - DOI:

<https://doi.org/10.23637/ERADOC-1-183>

63/C/3.1

LONG TERM LIMING EXPERIMENT - SPRING BEANS 1963

(LL and WLL)

Effect of lime on the yield and composition of crops and on the status of P and K in soils - Rothamsted (R) Sawyers I and Woburn (W) Stackyard Series C 1963, the second year.

Design (each field): 2 randomised blocks of 16 plots each.

Area of each plot (acres): 0.0289. Area harvested: 0.0121.

Treatments. All combinations of:-

Ground chalk (tons per acre): Sawyers I (R): None, 2, 4 applied in March 1962 (O, A and B), 8 (6 in March 1962, 2 in winter 1962 - 63 in divided dressings) (C). Stackyard Series C (W): None, 2 applied in Spring 1962 (O and A), 4.75 (4 in spring, 0.75 in October 1962) (B), 7.5 (6 in spring, 1.5 in October 1962) (C).

P: None, 0.5 cwt  $P_{2O_5}$  per acre as superphosphate (cumulative).

K: None, 1.0 cwt  $K_{2O}$  per acre as muriate of potash (cumulative).

The pH ranges between plots after harvest 1962 were as follows:-

Field	Chalk per acre (Spring 1962)	pH range
Sawyers I (R)	None	4.8 - 5.2
	2 tons	6.0 - 6.4
	4 tons	6.8 - 7.2
	6 tons	7.1 - 7.4
Stackyard Series C (W)	None	5.7 - 6.2
	2 tons	6.7 - 7.1
	4 tons	7.1 - 7.4
	6 tons	7.2 - 7.4

Cultivations, etc.

Sawyers I (R): Ground chalk applied at 1 ton per acre to 'C' plots: Dec 4, 1962. Ploughed: Mar 27, 1963. Ground chalk applied at 1 ton per acre to 'C' plots: Apr 1. Superphosphate and muriate of potash applied: Apr 3. Seed drilled at 200 lb per acre: Apr 8. Sprayed with simazine at 1 lb in 40 gallons per acre: Apr 18. Sprayed with demeton-methyl at 6 fluid oz in 40 gallons per acre: June 14. Combine harvested: Oct 18. Variety: Tick 30B. Previous crops: Potatoes and fallow 1960, potatoes and fallow 1961.

Stackyard Series C (W): Ploughed: Oct 12, 1962. Ground chalk applied at 0.75 tons per acre to 'B' plots and at 1.5 tons per acre to 'C' plots: Oct 19. Superphosphate and muriate of potash applied: Mar 13, 1963. Seed drilled at 200 lb per acre: Mar 27. Sprayed with simazine at 1 lb in 40 gallons per acre: Apr 8. Sprayed with demeton-methyl at 6 fluid oz in 40 gallons per acre: June 13. Combine harvested: Sept 21. Variety: Tick 30B. Previous crops: Barley 1960, sugar beet 1961.

63/C/3.2

Notes: (1) Samples were taken for counts of pods and beans.  
 (2) For details of the previous year's results see 'Numerical Results of the Field Experiments' 62/C/8.

Standard errors per plot. Grain (at 85% dry matter):  
 Sawyers I (R) 3.42 cwt per acre or 17.8% (15 d.f.)  
 Stackyard Series C(W) 2.37 cwt per acre or 15.1% (15 d.f.)

Summary of Results

Sawyers I (R)

Grain (at 85% dry matter): cwt per acre

	Ground chalk: tons per acre				Mean
	None	2	4	8	
Mean ( $\pm 1.21$ )	10.7	20.7	23.0	22.5	19.2
P <sub>20</sub> : cwt per acre	( $\pm 1.71$ )				
None	10.3	22.5	22.5	21.1	19.1
0.5	11.1	18.8	23.5	23.9	19.3
Diff. ( $\pm 2.42$ )	+0.8	-3.7	+1.0	+2.8	+0.2 ( $\pm 1.21$ )
K <sub>20</sub> : cwt per acre					
None	10.5	19.1	21.9	20.8	18.0
1.0	11.0	22.3	24.1	24.2	20.4
Diff. ( $\pm 2.42$ )	+0.5	+3.2	+2.2	+3.4	+2.4 ( $\pm 1.21$ )
	P <sub>20</sub> : cwt per acre				
	None	0.5			
	K <sub>20</sub> cwt per acre				
	( $\pm 1.21$ )				
None	18.4	17.7			
1.0	19.8	20.9			

Mean dry matter % as harvested: 68.7

63/C/3.3

Sawyers I (R)

Straw (at 85% dry matter): cwt per acre

	Ground chalk: tons per acre				Mean
	None	2	4	8	
Mean	7.6	14.1	17.3	14.8	13.5
<u>P205</u> : cwt per acre					
None	7.3	14.4	17.0	14.2	13.2
0.5	7.9	13.8	17.6	15.4	13.7
Diff.	+0.6	-0.6	+0.6	+1.2	+0.5
<u>K20</u> : cwt per acre					
None	7.4	13.1	15.4	11.8	11.9
1.0	7.8	15.1	19.2	17.9	15.0
Diff.	+0.4	+2.0	+3.8	+6.1	+3.1
	<u>P205</u> : cwt per acre	None	0.5		
<u>K20</u> : cwt per acre					
None	12.0	11.8			
1.0	14.4	15.6			

Mean dry matter % as harvested: 46.2



63/c/3.4

Stackyard Series C (W)

Grain (at 85% dry matter): cwt per acre

	Ground chalk: tons per acre				Mean
	None	2.00	4.75	7.50	
Mean ( $\pm 0.84$ )	12.4	17.5	16.5	16.5	15.7
P <sub>205</sub> : cwt per acre	( $\pm 1.18$ )				
None	11.2	15.8	17.0	15.4	14.9
0.5	13.7	19.2	16.0	17.7	16.6
Diff. ( $\pm 1.68$ )	+2.5	+3.4	-1.0	+2.3	+1.7 ( $\pm 0.84$ )
K <sub>20</sub> : cwt per acre					
None	11.6	14.3	14.7	15.2	13.9
1.0	13.3	20.7	18.3	17.9	17.5
Diff. ( $\pm 1.68$ )	+1.7	+6.4	+3.6	+2.7	+3.6 ( $\pm 0.84$ )
	P <sub>205</sub> : cwt per acre				
	None	0.5			
K <sub>20</sub> : cwt per acre	( $\pm 0.84$ )				
None	13.4	14.5			
1.0	16.4	18.7			

Mean dry matter % as harvested: 76.5