

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/AR/C/5 Methods of Application of Fertiliser 1962-63 - Potatoes

Rothamsted Research

Rothamsted Research (1964) *63/R/AR/C/5 Methods of Application of Fertiliser 1962-63 - Potatoes ; Yields Of The Field Experiments 1963*, pp 143 - 144 - DOI:

<https://doi.org/10.23637/ERADOC-1-183>

63/c/5.1

METHODS OF APPLICATION OF FERTILISER 1963 - 64.

(AR)

Methods of application of fertiliser - Great Knott II 1963, the first year - potatoes.

Design: 3 randomised blocks of 12 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0133 acres.

Treatments. None (0) (3 plots per block), and all combinations of:-
Levels of compound fertiliser (13% N, 13% P₂O₅, 20% K₂O) to supply (cwt per acre):-

N	P ₂ O ₅	K ₂ O	
0.66	0.66	1.02	(1)
1.32	1.32	2.03	(2)
2.00	2.00	3.07	(3)

Methods of application: Broadcast (B), placed (P), broadcast and rotary cultivated in (BR).

Note: The experiment is designed to include an additional factor applied to the 1964 wheat crop, viz. PK broadcast on seedbed and 'Nitro-Chalk' as spring top dressing to supply:-
N, P₂O₅ and K₂O at levels as (0), (1), (2) above.

Basal dressing: None.

Cultivations, etc.: Ploughed: Mar 28 - Apr 10, 1963. Rotary cultivated, potatoes planted: May 7. Earthed up: June 27. Sprayed with maneb at 1.2 lb in 20 gallons per acre: July 10. Sprayed with copper fungicide at 2.3 lb copper in 20 gallons per acre: July 29, and again at the same rate plus 0.35 pint menazon per acre: Aug 15. Sprayed with undiluted BOV at 16 gallons per acre: Sept 23. Lifted: Oct 21. Variety: King Edward. Previous crops: Barley 1961, spring beans 1962.

Standard error per plot.

Total tubers: 0.786 tons per acre or 7.3% (24 d.f.)

Erratum to 'Numerical Results of the Field Experiments' 1962 page 62/c/9.1, 'Total tubers: tons per acre'. Delete the standard error '(±0.181)' under the general mean.

63/C/5.2

Summary of Results

Method of application of fertiliser	Level of compound fertiliser				Mean	
	0	1	2	3		
<u>Total tubers: tons per acre</u>						
		(±0.454)			(±0.262)	
Broadcast		10.86	12.40	13.10	12.12	
Placed		10.08	11.93	13.83	11.95	
Broadcast and rotovated in		9.51	12.42	13.98	11.97	
Mean (±0.262)		7.23	10.15	12.25	13.64	10.81*

Percentage ware (1.5 inch riddle)

Broadcast		96.3	94.8	95.3	95.5	
Placed		93.4	94.1	93.7	93.7	
Broadcast and rotovated in		95.4	95.8	95.6	95.6	
Mean		93.8	95.1	94.9	94.9	94.6*

Level of compound fertiliser

	cwt per acre		
	N	P ₂ O ₅	K ₂ O
(0)		None	
(1)	0.66	0.66	1.02
(2)	1.32	1.32	2.03
(3)	2.00	2.00	3.07

* General mean