


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/AO/C/8 Decline of Take-all - Winter Wheat

Rothamsted Research

Rothamsted Research (1964) *63/R/AO/C/8 Decline of Take-all - Winter Wheat ; Yields Of The Field Experiments 1963*, pp 149 - 150 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/C/8.1

DECLINE OF TAKE-ALL

(AO)

The effect of crop sequences on the decline of take-all (Ophiobolus graminis) - Great Field I 1963, the first year.

Design: 3 randomised blocks of 6 plots each (5 of winter wheat, 1 of oats), using the plots of Series III of the Cereals and Beans Rotations Experiment (see 'Numerical Results of the Field Experiments' 61/C/1).

Area of each plot: 0.0305 acres. Area harvested: 0.0200 acres.

Treatments: Crop sequences:-

	1959	1960	1961	1962	1963	1964	1965	1966
1	W	W	WS	W	W	W	W	W
2	W	O	Be	W	W	W	W	W
3	WS	W	WS	W	O	W	W	W
4	B	W	B	W	W	W	O	W
5	W	O	WS	W	W	W	W	W
6	O	W	WS	W	W	O	W	W

O = Oats, Be = Spring beans, WS = Spring wheat, W = Winter wheat, B = Barley.

Basal dressings per acre: 2.5 cwt compound fertiliser (20% P₂O₅, 20% K₂O) combine drilled. 1.0 cwt N to wheat and 0.4 cwt N to oats applied in spring as 'Nitro-Chalk'.

Cultivations, etc.: Ploughed: Oct 29, 1962. Winter wheat drilled at 2.5 bushels per acre: Nov 14. 'Nitro-Chalk' applied to winter wheat: Mar 25, 1963. Oats drilled at 4 bushels per acre, 'Nitro-Chalk' applied to oats: Apr 9. Winter wheat sprayed with mecoprop/2,4-D at 7 pints in 40 gallons per acre: May 16. Oats sprayed with MCPA/dichlorprop at 3.2 pints in 40 gallons per acre: May 27. Combine harvested: Sept 10. Varieties: winter wheat - Cappelle, oats - Condor.

Notes: (1) Yields were only taken for winter wheat.
(2) Estimates were made on 5 occasions of the incidence of take-all on wheat.

Standard error per plot. Winter wheat.
Grain (at 85% dry matter): 1.91 cwt per acre or 7.3% (8 d.f.)

63/c/8.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Crop in	W	O	W	B	W	Mean
1959	W	W	O	W	O	
1960	WS	WS	WS	B	Be	
1961	W	W	W	W	W	
1962						
	28.1	27.0	24.4	25.9	25.5	26.2
			(±1.10)			

Mean dry matter % as harvested: 81.4