
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1963



Full Table of Content

63/R/HWF/A/3 Hoosfield - Wheat After Fallow

Rothamsted Research

Rothamsted Research (1964) 63/R/HWF/A/3 Hoosfield - Wheat After Fallow; Yields Of The Field Experiments 1963, pp 15 - 15 - DOI: https://doi.org/10.23637/ERADOC-1-183

63/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1963

(HWF)

Without manure 1851 and since

For history, treatments etc., see 'Details of the Classical and Long Term Experiments' 1956.

The cropped plots were subdivided and Squarehead's Master 13/4 was sown on the South side, Cappelle on the North.

Area of each plot (acres): Squarehead's Master 13/4 - 0.0690. Cappelle - 0.0552. Area harvested: 0.0365.

Cultivations, etc.:

Cropped plots. Ploughed: Oct 24, 1962. Seed drilled at 3 bushels per acre: Nov 23. Combine harvested: Sept 10.

Fallowed plots. Ploughed three times: Oct 24, 1962, May 31, 1963, July 16.

Note: Counts of straw number and estimates of eyespot (Cercosporella herpotrichoides) and take-all (Ophiobolus graminis) were made.

<u> </u>	Summary of Resu	ilts		
Plot	B3	B4	Bl	
No. of years of fallow	1	1	3	Mean
Variety Grain (at 8)	5% dry matter):	cwt per acr	re_	
Squarehead's Master 13/4 Cappelle	12.4	7.8 13.3	12.4	10.9
Mean	12.0	10.5	14.5	12.4
Straw (at 8	% dry matter):	cwt per acr	<u>'e</u>	
Squarehead's Master 13/4 Cappelle	17.4 16.2	9.5 15.9	16.7	14.5 15.3
Mean	16.8	12.7	15.2	14.9