


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1963

[Full Table of Content](#)



---

## 63/R/AQ/C/10 Cereal Disease Reference Plots - Wheat

### Rothamsted Research

Rothamsted Research (1964) *63/R/AQ/C/10 Cereal Disease Reference Plots - Wheat* ; Yields Of The Field Experiments 1963, pp 153 - 154 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/c/10.1

CEREAL DISEASE REFERENCE PLOTS

(AQ)

The effect of crop sequences on the incidence of cereal diseases - Pennell's Piece 1963, the first year.

Crops in 1963: Winter wheat, spring wheat, beans and oats.

Design: 4 randomised blocks of 4 plots each, 2 blocks for winter and 2 for spring wheat. In addition 1 plot per block was sown with oats and 1 with beans.

Area of each plot: 0.0180 acres. Area harvested: 0.0119 acres.

Treatments: Crop sequences:-

	1963	1964	1965	1966	1967
1	W	W	W	Be	O
2	W	W	Be	O	W
3	W	Be	O	W	W
4	Be	O	W	W	W
5	O	W	W	W	Be
6	W	W	W	W	W

W = Winter wheat (plots 1 - 12) and Spring wheat (plots 13 - 24),  
O = Oats, Be = Spring beans.

Basal dressings (cwt per acre):

N as 'Nitro-Chalk':- 1.0 N to winter wheat as spring top dressing.  
0.6 N to spring wheat in seedbed.  
0.4 N to oats in seedbed.

None to spring beans.

PK as compound fertiliser (14%  $P_2O_5$ , 28%  $K_2O$ ):-

0.5  $P_2O_5$ , 1.0  $K_2O$  to spring beans placement drilled.

0.3  $P_2O_5$ , 0.6  $K_2O$  to all other crops combine drilled.

Cultivations, etc.: Ploughed: Oct 18 - 26, 1962. Winter wheat:-  
Seed drilled at 2.5 bushels per acre: Oct 31, 1962. 'Nitro-Chalk'  
applied: Mar 13, 1963. Sprayed with mecoprop/2,4-D at 7 pints in  
40 gallons per acre: May 16. Combine harvested: Sept 10. Variety:  
Cappelle.

63/c/10.2

Spring wheat: Seed drilled at 3 bushels per acre: Apr 11, 1963.

'Nitro-Chalk' applied: Apr 18. Sprayed with methoxychlorobenzoic acid/MCPA (MBA/MCPA) at 4 pints in 40 gallons per acre: June 6. Combine harvested: Sept 12. Variety: Jufy I.

Oats: Seed drilled at 4 bushels per acre: Apr 9, 1963. 'Nitro-Chalk' applied: Apr 18. Sprayed with methoxychlorobenzoic acid/MCPA (MBA/MCPA) at 4 pints in 40 gallons per acre: June 6. Combine harvested: Sept 9. Variety: Condor.

Spring beans: Seed drilled at 200 lb per acre: Apr 9, 1963. Sprayed with demeton-methyl at 6 fluid oz in 40 gallons per acre: June 14. Combine harvested: Oct 3. Variety: Tick 30B.

Previous crops: Sugar beet 1961, spring wheat 1962.

Note: (1) Yields were only taken for winter and spring wheat.  
(2) Estimates of the incidence of take-all (Ophiobolus graminis), eyespot (Cercospora herpotrichoides) and sharp eyespot (Corticium solani) were made on 4 occasions for winter and 2 for spring wheat.

Standard errors per plot. Grain (at 85% dry matter):  
Winter wheat: 7.72 cwt per acre or 24.2% (6 d.f.)  
Spring wheat: 2.80 cwt per acre or 8.7% (6 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Winter wheat	Spring wheat
31.9	32.3
(±2.73)	(±0.99)

Mean dry matter % as harvested: Winter wheat 78.6  
Spring wheat 73.1