


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1963

[Full Table of Content](#)



---

### 63/W/WW101/DA/2 Winter Wheat - Varieties and N

#### Rothamsted Research

Rothamsted Research (1964) *63/W/WW101/DA/2 Winter Wheat - Varieties and N* ; Yields Of The Field Experiments 1963, pp 157 - 158 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/Da/2.1

WINTER WHEAT

(WW 101)

Levels and times of application of nitrogen - Woburn Butt Close 1963.

Design: 4 replicates of 20 treatments (4 x 5 factorial) arranged in 8 randomised blocks of 10 plots each, with one extra plot per block receiving no nitrogen other than basal.

Area of each plot: 0.0206 acres. Area harvested: 0.0137 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.5, 1.0, 1.5, 2.0 cwt N per acre as 'Nitro-Chalk'.

Times of application (intended): February\* (F), April (A), May (M), in two equal divided dressings (FM), in three (FAM).

\*See cultivations.

Basal dressing: 300 lb compound fertiliser (6% N, 15% P<sub>2</sub>O<sub>5</sub>, 15% K<sub>2</sub>O) per acre combine drilled.

Cultivations, etc.: Ploughed (after beans): Oct 10 1962, (after carrots): Oct 30. Seed drilled at 2.75 bushels per acre: Nov 1. 'Nitro-Chalk' applied: F - Mar 14, 1963, A - Apr 10, M - May 10. Sprayed with TBA/MCPA at 4 pints in 40 gallons per acre: May 9. Combine harvested: Sept 13. Variety: Cappelle. Previous crops: Plots 101 - 144, barley 1961, carrots 1962: plots 145 - 188, barley 1961, spring beans 1962.

Standard error per plot.

Grain (at 85% dry matter): 5.13 cwt per acre or 16.5% (53 d.f.)

63/Da/2.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	N: cwt per acre (in addition to basal)				Mean
	0.5	1.0	1.5	2.0	
Time of application	(±2.56)				(±1.28)
F	34.0	37.9	31.3	29.6	33.2
A	31.4	34.0	27.6	30.7	30.9
M	23.5	30.0	24.7	31.2	27.4
FM	27.8	37.3	35.0	26.9	31.8
FAM	25.5	35.1	34.5	32.2	31.8
Mean (±1.15)	28.4	34.9	30.6	30.1	31.0

Mean of plots receiving no nitrogen other than basal: 11.0

Mean dry matter % as harvested (all plots): 79.8

Time of application of N

F = Mar 14  
 A = Apr 10  
 M = May 10