


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1963

[Full Table of Content](#)



---

## 63/R/RW201/DA/3 Wheat - Varieties and N

### Rothamsted Research

Rothamsted Research (1964) *63/R/RW201/DA/3 Wheat - Varieties and N* ; Yields Of The Field Experiments 1963, pp 159 - 160 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/De/3.1

WHEAT

(RW 201)

Varieties and nitrogen - Long Hoos V 1963.

Design: 4 randomised blocks of 24 plots each, blocks being divided into 2 sub-blocks each, one of 15 winter wheat and one of 9 spring wheat plots.

Area of each plot: 0.0193 acres. Area harvested: 0.0129 acres.

Treatments: All combinations of:-

Varieties: Winter wheat:- Cappelle (Ca), Champlein (Ch),  
Prestige (Pr), Squarehead's Master (Sq).

Spring wheat:- July I (Ju), Opal (Op), Prestige (Pr).

Nitrogen: None (to winter wheat except Champlein), 0.5, 0.75,  
1.0 cwt N per acre as 'Nitro-Chalk'.

Basal dresssiinnngg: 2.5 cwt per acre compound fertiliser (20%  $P_{2}O_{5}$ ,  
20%  $K_{2}O$ ) combine drilled.

Cultivations, etc.: Ploughed: Nov 13, 1962. Winter wheat drilled at 2.75 bushels per acre: Dec 17. Spring wheat drilled at 3 bushels per acre: Apr 19, 1963. 'Nitro-Chalk' applied: Spring wheat - Apr 23, winter wheat - Apr 27. Sprayed with mecoprop/2,4-D: Winter wheat at 7 pints in 40 gallons per acre - May 16, spring wheat at 6 pints in 40 gallons per acre - May 24. Combine harvested: Winter wheat - Sept 10, spring wheat - Sept 13. Previous crops: Spring wheat 1961, potatoes 1962.

Standard errors per plot. Grain (at 85% dry matter):

Winter wheat 3.72 cwt per acre or 11.0% (39 d.f.)

Spring wheat 1.33 cwt per acre or 3.6% (24 d.f.)

63/Da/3.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Winter wheat

N: cwt per acre	Variety				Mean
	Ca	Ch	Pr	Sq	
	(±1.86)				(±0.93)
None	25.7		21.0	24.5	23.7 (1)
0.50	34.7	36.3	34.9	27.1	33.2
0.75	39.5	42.1	39.9	29.5	37.7
1.00	39.9	44.7	41.3	28.8	38.7
Mean (excl. none) (±1.07)	38.0	41.0	38.7	28.4	36.5*

(1) (±1.07)

\*Mean of plots receiving N. General mean 33.9

Mean dry matter % as harvested: 79.2

Spring wheat

N: cwt per acre	Variety			Mean
	Ju	Op	Pr	
	(±0.66)			(±0.38)
0.50	37.6	37.7	30.8	35.3
0.75	38.2	41.1	31.6	37.0
1.00	40.3	42.7	31.4	38.1
Mean (±0.38)	38.7	40.5	31.2	36.8

Mean dry matter % as harvested: 73.7

Varieties

Winter wheat

Ca = Cappelle, Ch = Champlain, Pr = Prestige,  
Sq = Squarehead's Master.

Spring wheat

Ju = July I, Op = Opal, Pr = Prestige